


## Functional regulation of estrogen receptor pathway by the dynein light chain 1

Suresh K Rayala, Petra den Hollander, Seetharaman Balasenthil, Zhibo Yang, Russell R. Broaddus & Rakesh Kumar

EMBO reports (2005) 6, 538–545. doi:10.1038/sj.embor.7400417

An incorrect version of Fig 1B in the article by Rayala *et al* was published in the June 2005 issue of *EMBO reports*. The entire figure is reprinted opposite in its correct form. The text of the article remains unchanged.

doi:10.1038/sj.embor.7400569



**Fig 1** | DLC1 is an E2-responsive gene. (A) DLC1 mRNA is increased after oestrogen stimulation. MCF-7 cells were treated with different concentrations of E2 for 16 h and analysed by northern blotting. (B) Effect of actinomycin D on the ability of E2 to upregulate DLC1 mRNA in MCF-7 cells. Cells were collected at different time points after treatment with E2 and/or actinomycin D (10 mg/ml) and DLC1 mRNA was analysed by northern blotting. (C) Upregulation of DLC1 promoter activity in MCF-7 and Ishikawa cells treated with E2 for 24 h. Cells were transfected with ERE-luc for 24 h, treated with E2 for 24 h and ERE-luc activity was measured. (D) Upregulation of DLC1 protein in MCF-7 and Ishikawa cells treated with E2 for 24 h.

## UVA inactivates protein tyrosine phosphatases by calpain-mediated degradation

Pawan Gulati, Boyka Markova, Martin Göttlicher, Frank-D. Böhmer & Peter A. Herrlich

EMBO reports (2004) 5, 812–817. doi:10.1038/sj.embor.7400190

Due to a calibration error, the UVA doses presented by P. Gulati *et al* in the August 2004 issue of *EMBO Reports* were 20.4-fold too low. The UVA lamp has been recalibrated with a new UV Lightmeter (UV-340, Lutron, Taiwan). Despite this correction, the

doses of UVA inducing the calpain-dependent cleavage of protein tyrosine phosphatases are still in the range of physiological exposure.

doi:10.1038/sj.embor.7400568