A CLASSIFICATION OF EQUIVARIANT PRINCIPAL BUNDLES OVER NONSINGULAR TORIC VARIETIES

INDRANIL BISWAS, ARIJIT DEY, AND MAINAK PODDAR

ABSTRACT. We classify holomorphic as well as algebraic torus equivariant principal G-bundles over a nonsingular toric variety X, where G is a complex linear algebraic group. It is shown that any such bundle over an affine, nonsingular toric variety admits a trivialization in equivariant sense. We also obtain some splitting results.

1. Introduction

Denote the algebraic torus $(\mathbb{C}^*)^n$ by T. Let N denote the group of 1-parameter subgroups of T; so N is isomorphic to \mathbb{Z}^n . Let M denote the group of characters of T. Then $M = Hom_{\mathbb{Z}}(N,\mathbb{Z})$ by restricting a character to the 1-parameter subgroups (the endomorphisms of \mathbb{C}^* is \mathbb{Z}).

Let X be a complex manifold (resp. smooth complex algebraic variety) equipped with a holomorphic (resp. algebraic) left-action of T, and let G be a complex linear algebraic group. A T-equivariant holomorphic (resp. algebraic) principal G-bundle on X is a holomorphic (resp. algebraic) principal G-bundle $\pi: \mathcal{E} \to X$ with T acting holomorphically (resp. algebraically) on \mathcal{E} , such that π is T-equivariant and

$$t(z \cdot g) = (tz) \cdot g \ \forall \ t \in T, \ g \in G, \ z \in \mathcal{E}$$

(this means that the actions of T and G on \mathcal{E} commute).

Now consider $X = X_{\Xi}$ to be a nonsingular complex toric variety of dimension n corresponding to a fan Ξ . Our main goal is to give a description of isomorphism classes of holomorphic (resp. algebraic) T-equivariant principal G-bundles over X; this is carried out in Theorem 3.2. This description is in terms of equivalence classes of data $\{\rho_{\sigma}, P(\tau, \sigma)\}$, where σ runs over maximal cones of Ξ , while $\rho_{\sigma}: T \to G$ is a holomorphic (equivalently algebraic) group homomorphism and $P(\tau, \sigma)$ is a G-valued 1-cocycle. This data $\{\rho_{\sigma}, P(\tau, \sigma)\}$ is required to satisfy certain conditions. In particular, the data depends on choice of projections $\pi_{\sigma}: T \to T_{\sigma}$ (see section 2). Here T_{σ} denotes the stabilizer of the orbit corresponding to the cone σ . However, if every maximal cone is of top dimension (such as for a complete toric variety), then such a choice of projections is not necessary.

Note that the description of isomorphism classes is the same in both the algebraic and holomorphic cases. We prove that every isomorphism class of holomorphic bundles contains an algebraic representative; see Remark 3.3.

²⁰¹⁰ Mathematics Subject Classification. 32L05, 14M25, 55R91.

Key words and phrases. Toric varieties, equivariant bundles, principal bundles.

As an application of the above description, we prove that if G is nilpotent then the principal G-bundle admits an equivariant reduction of structure group to a torus (meaning the bundle is split). For G abelian, this splitting result appeared in [1].

In section 4, we give a more combinatorial description of the isomorphism classes of T-equivariant principal G-bundles in terms of characters of T and G-valued 1-cocycles; see Theorems 4.5 and 4.6. These generalize Kaneyama's description [7] of the isomorphism classes of T-equivariant vector bundles over complete nonsingular toric varieties in two directions. Firstly, the assumption of completeness on the base is removed. Secondly, the bundles are generalized from vector to principal bundles.

If G has a normal maximal torus, then a T-equivariant principal G-bundle over X has an equivariant reduction of structure group to the maximal torus; see Lemma 4.3. Our treatment also makes clearer the approach of Kaneyama. This helps us unearth a gap in the proof of the splitting of small rank equivariant vector bundles over projective space in [8]; see Remark 4.7.

In Lemma 2.11 we show that the automorphism group of a T-equivariant holomorphic (or algebraic) principal G-bundle over a toric variety is a subgroup of G that contains the center of G.

A crucial tool used by us is the following equivariant Oka principle due to Heinzner and Kutzschebauch [5]. The algebraic analogue of this is derived in the toric case in Lemma 2.7.

Theorem 1.1. [5] Suppose K is any complex reductive group acting linearly on \mathbb{C}^k . Then any K-equivariant holomorphic principal G-bundle \mathcal{E} over \mathbb{C}^k is equivariantly isomorphic to $\mathbb{C}^k \times \mathcal{E}(0)$ where the latter has diagonal K-action.

Some of the basic lemmas regarding local action functions in section 2 have appeared before in [1]. We include them here for the convenience of the reader.

In the topological case, the classification of $(S^1)^n$ -equivariant principal G-bundles over a toric manifold, where G is a compact Lie group, is contained in the results of Hambleton and Hausmann [3]. The classification of $(\mathbb{C}^*)^n$ -equivariant topological principal bundles over a toric manifold appears to be an interesting open problem.

2. Distinguished sections

Denote by $\Xi(d)$ the set of all d-dimensional cones in Ξ . For a cone σ in Ξ , denote the corresponding affine variety and T-orbit by X_{σ} and O_{σ} respectively. Let T_{σ} denote the stabilizer of any point in O_{σ} ; it is independent of the point because T is abelian. We recall that each O_{σ} may be given a group structure using a specific identification with T/T_{σ} , see [2, p. 53] or [10, Proposition 1.6]. The principal orbit O may be identified with T.

For each σ , fix once and for all, an embedding $\iota_{\sigma}: O_{\sigma} \hookrightarrow T$ that splits the exact sequence

$$1 \to T_{\sigma} \to T \to O_{\sigma} \to 1$$
.

As we are in the nonsingular case, such an embedding may be obtained by extending a set of independent, primitive generators of σ to a basis of the lattice N. A splitting,

as above, induces a direct product decomposition $T = T_{\sigma} \times \iota_{\sigma}(O_{\sigma})$. Let $\pi_{\sigma} : T \to T_{\sigma}$ be the projection associated to this decomposition of T.

Fix any $\sigma \in \Xi$. Let $X = X_{\sigma}$. There exists an affine toric variety A_{σ} with the dense orbit isomorphic to T_{σ} , such that X is T-equivariantly isomorphic to $A_{\sigma} \times O_{\sigma}$. Under this isomorphism, the orbit O_{σ} in X is identified with $\{p\} \times O_{\sigma}$, where p is the T_{σ} -fixed point of A_{σ} . Moreover, there exist isomorphisms $a_{\sigma} : T_{\sigma} \to (\mathbb{C}^*)^{\dim(\sigma)}$ and $\phi_{\sigma} : A_{\sigma} \to \mathbb{C}^{\dim(\sigma)}$, satisfying

$$\phi_{\sigma}(tx) = a_{\sigma}(t)\phi_{\sigma}(x)$$

for all $x \in A_{\sigma}$ and $t \in T_{\sigma}$.

Let G denote a complex linear algebraic group. Suppose \mathcal{E} is a T-equivariant holomorphic (resp. algebraic) principal G-bundle over X which admits a holomorphic (resp. algebraic) trivialization. Let $s: X \to \mathcal{E}$ be any holomorphic (resp. algebraic) section. We encode the T-action on \mathcal{E} as follows:

Definition 2.1. For any $x \in X$ and $t \in T$, define $\rho_s(x,t) \in G$ by

$$ts(x) = s(tx) \cdot \rho_s(x,t)$$
.

Since the action of G on each fiber of \mathcal{E} is free and transitive, it follows that $\rho_s(x,t)$ is well-defined and it is holomorphic (resp. algebraic) in both x and t. We say that $\rho_s: X \times T \to G$ is the local action function associated to s.

If $s'(x) = s(x) \cdot g(x)$ is another holomorphic (resp. algebraic) section of \mathcal{E} , then it is straight-forward to check that

$$\rho_{s'}(x,t) = g(tx)^{-1}\rho_s(x,t)g(x).$$
(2.1)

Lemma 2.2. For any $t_1, t_2 \in T$, the equality

$$\rho_s(x, t_1 t_2) = \rho_s(t_2 x, t_1) \rho_s(x, t_2)$$

holds.

Proof. By the definition of ρ_s , we have $t_1t_2s(x) = s(t_1t_2x) \cdot \rho_s(x,t_1t_2)$. On the other hand,

$$t_1 t_2 s(x) = t_1(s(t_2 x) \cdot \rho_s(x, t_2)) = (t_1 s(t_2 x)) \cdot \rho_s(x, t_2) = s(t_1 t_2 x) \cdot \rho_s(t_2 x, t_1) \rho_s(x, t_2)$$
. The lemma follows from these.

Suppose δ is any subcone of σ . Then Lemma 2.2 implies that for any $x_{\delta} \in O_{\delta}$, the restriction

$$\rho_s(x_\delta,\cdot):T_\delta\to G$$

is a group homomorphism, where T_{δ} as before is the stabilizer.

Lemma 2.3. If $\rho_s(x,\cdot)$ is independent of x, then it defines a group homomorphism

$$\rho_{\mathfrak{s}}: T \to G$$
.

Conversely if $\rho_s(x_0,\cdot)$ is a group homomorphism for some $x_0 \in O$, then $\rho_s(x,\cdot)$ is independent of x.

Proof. The first part follows immediately from Lemma 2.2.

To prove the second part, assume that $\rho_s(x_0,\cdot)$ is a group homomorphism for some $x_0 \in O$. We have

$$\rho_s(x_0, t)\rho_s(x_0, u) = \rho_s(x_0, tu) = \rho_s(ux_0, t)\rho_s(x_0, u)$$

for all $u, t \in T$. Therefore, for any $u \in T$, we have

$$\rho_s(ux_0,\cdot) = \rho_s(x_0,\cdot).$$

Then, since ρ_s is continuous, the second part of the lemma follows from the denseness of O in X.

Lemma 2.4. Let X_1 and X_2 be affine toric varieties. Let $\alpha: X_1 \to X_2$ be an isomorphism of T-spaces up to an automorphism $a: T \to T$, meaning $\alpha \circ t = a(t) \circ \alpha$. Suppose $\pi_i: \mathcal{E}_i \to X_i$ is a T-equivariant trivial principal G-bundle for i = 1, 2. Let $\phi: \mathcal{E}_1 \to \mathcal{E}_2$ be an isomorphism of T-equivariant principal G-bundles over X, which is compatible with α and a:

$$\pi_2 \circ \phi = \alpha \circ \pi_1$$
 and $\phi \circ t = a(t) \circ \phi$.

Let s_1 be any section of \mathcal{E}_1 . Let s_2 be the section of \mathcal{E}_2 defined by $s_2(\alpha(x)) = \phi(s_1(x))$ for any $x \in X_1$. Then $\rho_{s_1}(x,t) = \rho_{s_2}(\alpha(x),a(t))$ for every $x \in X_1$, $t \in T$. In particular, if α and a are both identity, then $\rho_{s_1} = \rho_{s_2}$.

Proof. This follows from the following calculation:

$$s_{2}(a(t)\alpha(x)) \cdot \rho_{s_{2}}(\alpha(x), a(t))$$

$$= a(t)s_{2}(\alpha(x)) = a(t)\phi(s_{1}(x))$$

$$= \phi(ts_{1}(x)) = \phi(s_{1}(tx) \cdot \rho_{s_{1}}(x, t))$$

$$= \phi(s_{1}(tx)) \cdot \rho_{s_{1}}(x, t) = s_{2}(\alpha(tx)) \cdot \rho_{s_{1}}(x, t)$$

$$= s_{2}(a(t)\alpha(x)) \cdot \rho_{s_{1}}(x, t).$$

Definition 2.5. We say that a section s of \mathcal{E} is distinguished if

- $\rho_s(x,\cdot)$ is independent of x, and
- $\rho_s(x,\cdot)$ factors through the projection $\pi_\sigma: T \to T_\sigma$.

Lemma 2.6. If s is a distinguished section, then so is $s \cdot g$ for every $g \in G$.

Proof. The assertion follows from the fact that the actions of T and G commute. Note that $\rho_{s,q}(x,t) = g^{-1}\rho_s(x,t)g$ by (2.1).

Lemma 2.7. Consider any linear action of the torus $K = (\mathbb{C}^*)^k$ on \mathbb{C}^k . Suppose \mathcal{E} is a K-equivariant algebraic principal G-bundle over \mathbb{C}^k . Then \mathcal{E} is K-equivariantly isomorphic to $\mathbb{C}^k \times \mathcal{E}(0)$, where the latter has diagonal K-action.

Proof. In particular, \mathcal{E} is a K-equivariant holomorphic principal G bundle. Therefore, by Theorem 1.1 \mathcal{E} admits a K-equivariant holomorphic trivialization, $\mathbb{C}^k \times \mathcal{E}(0)$. Let s be a holomorphic section of \mathcal{E} , which corresponds to a constant section of the

trivialization $\mathbb{C}^k \times \mathcal{E}(0)$. Let $\rho_s : T \to G$ be the local action homomorphism corresponding to s. Let x_0 be any point in $(\mathbb{C}^*)^k$. Then

$$s(tx_0) = ts(x_0) \cdot \rho_s(t)^{-1}.$$

Since the homomorphism ρ_s and the T-action are algebraic, the holomorphic section s is algebraic over $(\mathbb{C}^*)^k$. This implies that s is in fact algebraic over \mathbb{C}^k . Thus \mathcal{E} admits an algebraic K-equivariant trivialization.

Lemma 2.8. Let \mathcal{E} be a T-equivariant holomorphic (resp. algebraic) principal G-bundle on a nonsingular affine toric variety X_{σ} . Then \mathcal{E} is trivial and it admits a distinguished section.

Proof. Recall that $X_{\sigma} = A_{\sigma} \times O_{\sigma}$, where A_{σ} is isomorphic to $\mathbb{C}^{\dim \sigma}$ and T_{σ} acts linearly on A_{σ} . Then by Theorem 1.1 (resp. Lemma 2.7), the restriction $\mathcal{E}|_{A_{\sigma}}$ is T_{σ} -equivariantly isomorphic to $A_{\sigma} \times \mathcal{E}(p)$, where p is the T_{σ} -fixed point of A_{σ} . Here the T_{σ} -action on the product $A_{\sigma} \times \mathcal{E}(p)$ is the natural diagonal action. Take any $e \in \mathcal{E}(p)$. Let $\rho: T_{\sigma} \to G$ be the homomorphism induced by the action of T_{σ} on $\mathcal{E}(p)$ defined by $tz = z \cdot \rho(t)$.

Let s be the section of $\mathcal{E}|_{A_{\sigma}}$ corresponding to the constant section e of $A_{\sigma} \times \mathcal{E}(p)$. Then $\rho_s = \rho$ by the definition of s.

Let y_0 be the identity element in O_{σ} . We may identify A_{σ} with $A_{\sigma} \times \{y_0\}$. Now extend s to a section of \mathcal{E} over X_{σ} by setting

$$s(x, ky_0) = \iota_{\sigma}(k)s(x, y_0) \tag{2.2}$$

for every $x \in A_{\sigma}$ and $k \in O_{\sigma}$. This shows that \mathcal{E} is trivial over X_{σ} .

The local action function of the section s over $A_{\sigma} \times \{y_0\}$ satisfies

$$\rho_s((x, y_0), h) = \rho_s((p, y_0), h) = \rho(h)$$

for all $h \in T_{\sigma}$. Since

$$hs(x, ky_0) = h\iota_{\sigma}(k)s(x, y_0) = \iota_{\sigma}(k)hs(x, y_0) = \iota_{\sigma}(k)s(hx, y_0) \cdot \rho_s((x, y_0), h) = s(hx, ky_0) \cdot \rho_s((x, y_0), h),$$

we deduce that

$$\rho_s((x, ky_0), h) = \rho_s((x, y_0), h) \tag{2.3}$$

for every $k \in O_{\sigma}$ and every $h \in T_{\sigma}$.

It follows easily from (2.2) that $\iota_{\sigma}(k)s(x,y) = s(x,ky)$ for any $y \in O_{\sigma}$. Then, using (2.3), we have

$$h\iota_{\sigma}(k)s(x,y) = hs(x,ky) = s(hx,ky) \cdot \rho_s((x,ky),h)$$

= $s(hx,ky) \cdot \rho_s((x,y_0),h) = s(hx,ky) \cdot \rho_s((p,y_0),h)$

for every $(h, \iota_{\sigma}(k)) \in T$ and $(x, y) \in A_{\sigma} \times O_{\sigma} = X_{\sigma}$. Therefore,

$$\rho_s((x,y),h\iota_{\sigma}(k)) = \rho_s((p,y_0),h) = \rho(h),$$

and the lemma follows.

Lemma 2.9. Let σ be a cone. Then the homomorphisms $\rho_s: T \to G$, induced by different distinguished sections s of $\mathcal{E}|_{X_{\sigma}}$, are equal up to conjugation by elements of G.

Proof. Take a distinguished section s. Let $x_{\sigma} = (p, y_0)$ be the fixed point of T_{σ} in X_{σ} corresponding to the identity element of O_{σ} . Then for any $x \in X_{\sigma}$, $\rho_s(x,t) = \rho_s(x_{\sigma},t)$. Note that

$$\rho_s(x,t) = \rho_s(x_{\sigma},t) = \rho_s(x_{\sigma},\pi_{\sigma}(t)),$$

since s is a distinguished section. Moreover,

$$\pi_{\sigma}(t)s(x_{\sigma}) = s(x_{\sigma}) \cdot \rho_{s}(x_{\sigma}, \pi_{\sigma}(t)).$$

Therefore, ρ_s is completely determined by $s(x_\sigma)$ and the canonical T_σ action on $\mathcal{E}(x_\sigma)$. If s' is another distinguished section, then there exists $g \in G$ such that $s'(x_\sigma) = s(x_\sigma) \cdot g$. Then $\rho_{s'} = g^{-1}\rho_s g$ follows by applying equation (2.1) with $x = x_\sigma$, $t \in T_\sigma$, and $g(x_\sigma) = g$. Note that by Lemma 2.6 all elements of G appear this way for distinguished sections.

Definition 2.10. For notational convenience, denote by ρ_{σ} a homomorphism from T to G induced by a distinguished section of $\mathcal{E}|_{X_{\sigma}}$. By Lemma 2.9, the homomorphism ρ_{σ} is unique up to conjugation by elements of G.

Lemma 2.11. Let \mathcal{E} be a T-equivariant holomorphic (resp. algebraic) principal G-bundle over a nonsingular toric variety X (not necessarily affine). Let $\operatorname{Aut}_{T}(\mathcal{E})$ be the group of T-equivariant automorphisms of \mathcal{E} that project to the identity map on X. Then $\operatorname{Aut}_{T}(\mathcal{E})$ is a subgroup of G that contains the center Z(G) of G.

Proof. Fix a point $x_0 \in O \subset X$ and also an element $e \in \mathcal{E}(x_0)$. Consider the map

$$\alpha: \operatorname{Aut}_{\tau}(\mathcal{E}) \to G$$

uniquely determined by the equation $\Phi(e) = e \cdot \alpha(\Phi)$. Note that given any $\Phi \in \operatorname{Aut}_T(\mathcal{E})$, by its T-equivariance and continuity, Φ is determined by the restriction $\Phi|_{\mathcal{E}(x_0)}$, while $\Phi|_{\mathcal{E}(x_0)}$ in turn is determined by $\Phi(e)$ using G-equivariance. Therefore the above map α is injective. For $\Phi_1 \in \operatorname{Aut}_T(\mathcal{E})$, we have

$$\Phi_2 \circ \Phi_1(e) = \Phi_2(e \cdot \alpha(\Phi_1)) = \Phi_2(e) \cdot \alpha(\Phi_1) = e \cdot \alpha(\Phi_2)\alpha(\Phi_1).$$

This proves that α is a group homomorphism.

Now take any $g \in Z(G)$. Choose a distinguished section s of \mathcal{E}_{σ} such that $s(x_0) = e$. Define $\Phi(e) = e \cdot g$. Then Φ determines an automorphism of $\mathcal{E}|_{O}$ using T and G equivariance as follows: $\Phi(te \cdot h) = te \cdot gh$. We need to check that Φ extends to the boundary strata of X, which consist of lower dimensional T-orbits. Note that

$$\Phi(te) = \Phi(ts(x_0)) = ts(x_0) \cdot g = s(tx_0) \cdot \rho_s(t)g.$$

On the other hand,

$$\Phi(te) \, = \, \Phi(ts(x_0)) \, = \, \Phi(s(tx_0) \cdot \rho_s(t)) \, = \, \Phi(s(tx_0)) \cdot \rho_s(t) \, .$$

Therefore,

$$\Phi(s(tx_0)) = s(tx_0) \cdot \rho_s(t)g\rho_s(t)^{-1} = s(tx_0) \cdot g$$

as $g \in Z(G)$. Any point x in a boundary stratum can be considered as a limit

$$\lim_{z\to 0}\lambda(z)x_0\,,$$

where $\lambda(z)$ is an algebraic curve in T. Now since s and the G-action are holomorphic (resp. algebraic), $\Phi(s(x)) = s(x) \cdot g$ defines an analytic (resp. algebraic) extension of Φ over X.

3. Isomorphism classes and admissible collections

Let X be a nonsingular toric variety of dimension n corresponding to a fan Ξ . Let \mathcal{E} be a holomorphic (resp. algebraic) T-equivariant principal G-bundle over X. Define $\mathcal{E}_{\sigma} := \mathcal{E}|_{X_{\sigma}}$ for any cone $\sigma \in \Xi$.

Let σ be a maximal cone in Ξ , and let s_{σ} be a distinguished section of \mathcal{E}_{σ} . Let $\rho_{\sigma}: T \to G$ be the corresponding homomorphism. The group T acts on $X_{\sigma} \times G$ as follows:

$$t(x,h) = (tx, \rho_{\sigma}(t)h). \tag{3.1}$$

Let $\psi_{\sigma}: \mathcal{E}_{\sigma} \to X_{\sigma} \times G$ be the trivialization of \mathcal{E}_{σ} defined by

$$\psi_{\sigma}(s_{\sigma}(x) \cdot h) = (x, h). \tag{3.2}$$

Note that

$$\psi_{\sigma}(ts_{\sigma}(x) \cdot h) = \psi_{\sigma}(s_{\sigma}(tx) \cdot \rho_{\sigma}(t)h) = (tx, \rho_{\sigma}(t)h). \tag{3.3}$$

Hence by (3.1) and (3.3) we get that ψ_{σ} in (3.2) is a *T*-equivariant trivialization.

Let σ, τ be any two maximal cones. Define $\phi_{\tau\sigma}: X_{\sigma} \cap X_{\tau} \to G$ as follows:

$$s_{\sigma}(x) = s_{\tau}(x) \cdot \phi_{\tau\sigma}(x). \tag{3.4}$$

The maps $\phi_{\tau\sigma}$'s are transition functions for \mathcal{E} as they satisfy

$$\psi_{\tau}\psi_{\sigma}^{-1}(x,h) = (x,\phi_{\tau\sigma}(x)h).$$
 (3.5)

By the T-equivariance property of \mathcal{E} we have

$$t(\psi_{\tau}\psi_{\sigma}^{-1}(x,h)) = \psi_{\tau}\psi_{\sigma}^{-1}(t(x,h)). \tag{3.6}$$

Using (3.1) and (3.5), from (3.6) it follows that

$$(tx, \rho_{\tau}(t)\phi_{\tau\sigma}(x)h) = (tx, \phi_{\tau\sigma}(tx)\rho_{\sigma}(t)h).$$

Therefore, we have $\phi_{\tau\sigma}(tx) = \rho_{\tau}(t)\phi_{\tau\sigma}(x)\rho_{\sigma}(t)^{-1}$.

Fix a point x_0 in the principal orbit O. Denote $\phi_{\tau\sigma}(x_0)$ by $P(\tau,\sigma)$. Note that $\rho_{\tau}(t)P(\tau,\sigma)\rho_{\sigma}(t)^{-1}$ extends to a holomorphic (resp. algebraic) function, namely $\phi_{\tau\sigma}$, on $X_{\tau\cap\sigma}$. Moreover, since $\phi_{\tau\sigma}$ are transition functions, $\{P(\tau,\sigma)\}$ satisfy the cocycle condition

$$P(\tau, \sigma)P(\sigma, \delta)P(\delta, \tau) = 1_G.$$

Finally, if we change $\{s_{\sigma}\}$ to $\{s_{\sigma} \cdot g_{\sigma}\}$, then $\{\rho_{\sigma}\}$ transforms to $\{g_{\sigma}^{-1}\rho_{\sigma}g_{\sigma}\}$, while $\{P(\tau,\sigma)\}$ transforms to $\{g_{\tau}^{-1}P(\tau,\sigma)g_{\sigma}\}$.

Definition 3.1. Let Ξ^* denote the set of maximal cones in Ξ . An admissible collection $\{\rho_{\sigma}, P(\tau, \sigma)\}$ consists of a collection of homomorphisms $\{\rho_{\sigma}: T \to G \mid \sigma \in \Xi^*\}$ and a collection of elements $\{P(\tau, \sigma) \in G \mid \tau, \sigma \in \Xi^*\}$ satisfying the following conditions:

(1) ρ_{σ} factors through $\pi_{\sigma}: T \to T_{\sigma}$.

- (2) For every pair (τ, σ) of maximal cones, $\rho_{\tau} P(\tau, \sigma) \rho_{\sigma}^{-1}$ extends to a G-valued holomorphic (equivalently regular algebraic) function over $X_{\sigma} \cap X_{\tau}$.
- (3) $P(\sigma, \sigma) = 1_G$ for all σ .
- (4) For every triple (τ, σ, δ) of maximal cones having nonempty intersection, the cocycle condition $P(\tau, \sigma)P(\sigma, \delta)$ $P(\delta, \tau) = 1_G$ holds.

Two such admissible collections $\{\rho_{\sigma}, P(\tau, \sigma)\}\$ and $\{\rho'_{\sigma}, P'(\tau, \sigma)\}\$ are equivalent if the following hold:

- (i) For every σ there exists g_σ ∈ G such that ρ'_σ = g_σ⁻¹ρ_σg_σ.
 (ii) For every pair (τ, σ), P'(τ, σ) = g_τ⁻¹P(τ, σ)g_σ, where g_σ and g_τ are as in (i) above.

Theorem 3.2. Let X be a nonsingular toric variety and G a complex linear algebraic group. Then the isomorphism classes of holomorphic (or algebraic) T-equivariant principal G-bundles on X are in one-to-one correspondence with equivalence classes of admissible collections $\{\rho_{\sigma}, P(\tau, \sigma)\}.$

Proof. Let \mathcal{E}_1 and \mathcal{E}_2 be two T-equivariant principal G-bundles on X which are Tequivariantly isomorphic. Fix such an isomorphism $\Phi: \mathcal{E}_1 \to \mathcal{E}_2$. Let $\{s^1_\sigma\}$ be a collection of distinguished sections which associates a class of admissible collections $\{\rho_{s^1_{\sigma}}, P^1(\tau, \sigma)\}\$ to $\stackrel{\smile}{\mathcal{E}}_1$. Let $s^2_{\sigma} = \Phi \circ s^1_{\sigma}$. Note that $\rho_{s^1_{\sigma}}(x, t) = \rho_{s^2_{\sigma}}(x, t)$ by Lemma 2.4. Hence the collection $\{s_{\sigma}^2\}$ is also distinguished. Then by (3.4) and the Gequivariance property of Φ we get that

$$s_{\sigma}^{2}(x_{0}) = \Phi(s_{\sigma}^{1}(x_{0})) = \Phi(s_{\tau}^{1}(x_{0})) \cdot \phi_{\tau\sigma}(x_{0}) = s_{\tau}^{2}(x_{0}) \cdot \phi_{\tau\sigma}(x_{0}).$$

Consequently, $P^2(\tau, \sigma) = P^1(\tau, \sigma)$. Therefore the admissible collection for \mathcal{E}_1 (with respect to $\{s^1_\sigma\}$) coincides with admissible collection for \mathcal{E}_2 (with respect to $\{s^2_\sigma\}$). However, for a different choice of collection of distinguished sections we get an equivalent admissible collection.

Thus we may associate an equivalence class of admissible collections to an isomorphism class of equivariant principal bundles. We will denote this assignment by

Conversely given an admissible collection $\{\rho, P\} = \{\rho_{\sigma}, P(\tau, \sigma)\}\$, we construct a principal G-bundle $E(\{\rho,P\})$ as follows. Let $\phi_{\tau\sigma}: X_{\sigma} \cap X_{\tau} \to G$ denote the holomorphic (equivalently regular algebraic) extension of $\rho_{\tau}P(\tau,\sigma)\rho_{\sigma}^{-1}$. Note that $\{\phi_{\tau\sigma}\}$ satisfies the cocycle condition. Therefore we may construct an algebraic (hence, holomorphic) principal G-bundle $E(\{\rho, P\})$ over X with $\{\phi_{\tau\sigma}\}$ as transition functions:

$$E(\{\rho, P\}) = (\bigsqcup_{\sigma} X_{\sigma} \times G) / \sim$$

where $(x,g) \sim (y,h)$ for $(x,g) \in X_{\sigma} \times G$ and $(y,h) \in X_{\tau} \times G$ if and only if

$$x = y, \quad x \in X_{\sigma} \cap X_{\tau} \text{ and } h = \phi_{\tau\sigma}(x)g.$$
 (3.7)

The G-action on each $X_{\sigma} \times G$ is defined by right multiplication. It is easy to check that these actions on different charts are compatible.

Define T action on each $X_{\sigma} \times G$ by $t(x,g) = (tx, \rho_{\sigma}(t)g)$. Note that if $(y,h) \in X_{\tau} \times G$ is equivalent to $(x,g) \in X_{\sigma} \times G$, then

$$t(y,h) = t(x,\phi_{\tau\sigma}(x)g) = (tx,\rho_{\tau}(t)\phi_{\tau\sigma}(x)g). \tag{3.8}$$

Now if x belongs to the open orbit $O = T \subset X_{\sigma} \cap X_{\tau}$, then

$$\phi_{\tau\sigma}(tx)\rho_{\sigma}(t) = \rho_{\tau}(tx)P(\tau,\sigma)\rho_{\sigma}(tx)^{-1}\rho_{\sigma}(t) = \rho_{\tau}(t)\rho_{\tau}(x)P(\tau,\sigma)\rho_{\sigma}(x)^{-1} = \rho_{\tau}(t)\phi_{\tau\sigma}(x).$$

Since both $\phi_{\tau\sigma}(tx)\rho_{\sigma}(t)$ and $\rho_{\tau}(t)\phi_{\tau\sigma}(x)$ are continuous in x on $X_{\sigma}\cap X_{\tau}$ and O is dense in $X_{\sigma}\cap X_{\tau}$,

$$\phi_{\tau\sigma}(tx)\rho_{\sigma}(t) = \rho_{\tau}(t)\phi_{\tau\sigma}(x) \quad \forall \ x \in X_{\sigma} \cap X_{\tau}. \tag{3.9}$$

From (3.8) and (3.9), we have

$$t(y,h) = (tx, \phi_{\tau\sigma}(tx)\rho_{\sigma}(t)g)$$

whenever $(x,g) \sim (y,h)$. Since $t(x,g) = (tx,\rho_{\sigma}(t)g)$, using (3.7) it follows that $t(y,h) \sim t(x,g)$ whenever $(x,g) \sim (y,h)$. In other words, the *T*-actions on the $X_{\sigma} \times G$ are compatible and define an (algebraic) action of *T* on $E(\{\rho,P\})$.

It is easy to check that $\mathbf{A}([E(\{\rho,P\})]) = [\{\rho,P\}]$. This shows that \mathbf{A} is surjective. Now we will show that \mathbf{A} is injective. Given a principal G-bundle \mathcal{E} on X, consider any representative $\{\rho,P\}$ of $\mathbf{A}([\mathcal{E}])$. We will construct an isomorphism $\Phi: \mathcal{E} \to E(\{\rho,P\})$. For this first note that $\{\rho,P\} = \{\rho_{\sigma},P(\tau,\sigma)\}$ is associated to a collection of distinguished sections $\{s_{\sigma}\}$. Given $e \in \mathcal{E}$, suppose $e \in \mathcal{E}_{\sigma}$. Then there exists a unique $g_e^{\sigma} \in G$ such that $e = s_{\sigma}(\pi(e)) \cdot g_e^{\sigma}$. Define

$$\Phi(e) = [(\pi(e), g_e^{\sigma})],$$

where the right hand side is the equivalence class of $(\pi(e), g_e^{\sigma})$ under the relation \sim defined in (3.7).

Using (3.4) it is easy to check that Φ is well-defined. Since $e \cdot h = s_{\sigma}(\pi(e)) \cdot g_e^{\sigma} h$, we have

$$\Phi(e \cdot h) = [\pi(e), g_e^{\sigma} h] = \Phi(e) \cdot h.$$

Therefore, Φ is a morphism of principal G-bundles. Since every morphism of principal G-bundles over a fixed base is an isomorphism, it follows that Φ is an isomorphism. Note that

$$te = ts_{\sigma}(\pi(e)) \cdot g_e^{\sigma} = s_{\sigma}(t\pi(e)) \cdot \rho_{\sigma}(t)g_e^{\sigma}.$$

Therefore,

$$\Phi(te) = [(t\pi(e), \rho_{\sigma}(t)g_{e}^{\sigma})] = t[\phi(e), g_{e}^{\sigma}] = t\Phi(e)$$
.

Hence Φ is T-equivariant. The inverse of Φ is then automatically T-equivariant.

Now suppose \mathcal{E}_1 , \mathcal{E}_2 are two equivariant principal G-bundle such that $\mathbf{A}[\mathcal{E}_1] = \mathbf{A}[\mathcal{E}_2] = [\{\rho, P\}]$. Then it follows from the above that

$$\mathcal{E}_1 \cong E(\{\rho, P\}) \cong \mathcal{E}_2$$
.

Therefore **A** is injective. This concludes the proof.

Remark 3.3. It follows from the proof of Theorem 3.2 that every isomorphism class of T-equivariant holomorphic principal G-bundles over X has an algebraic representative.

The following generalizes a similar result for abelian structure groups in [1].

Corollary 3.4. If G is a nilpotent group, then a T-equivariant holomorphic (or algebraic) principal G-bundle over a nonsingular toric variety admits an equivariant reduction of structure group to a maximal torus of G. In particular, if G is unipotent then the bundle is trivial with trivial T-action.

Proof. If G is nilpotent, then it is a direct product $K \times U$, where K is a maximal torus in G and U is unipotent. Homomorphisms from T to U are trivial. So the image of any homomorphism $\rho: T \to G$ lies in K.

Recall that if s_{σ} is a distinguished section, so is $s_{\sigma} \cdot g$ for any $g \in G$. Using this we can choose distinguished sections so that $P(\tau, \sigma) = 1_G$ for all pairs (τ, σ) of maximal cones. Then the transitions maps $\phi_{\tau\sigma} = \rho_{\tau}\rho_{\sigma}^{-1}$ take values in K. This implies the principal bundle has a reduction to K.

The claim about unipotent G follows easily from the fact that homomorphisms from T to G are trivial.

Example 3.5. Let (n_1, \ldots, n_k) be a partition of r. Let G be the nilpotent subgroup of $GL(r, \mathbb{C})$ consisting of all block diagonal matrices with k blocks of sizes n_1, \ldots, n_k respectively, where each block is upper triangular and the diagonal elements within a block are the same. Assume X is a complete, nonsingular toric variety with fan Ξ . Then the isomorphism classes of T-equivariant holomorphic (equivalently algebraic) principal G-bundles on X are parametrized by \mathbb{Z}^{dk} , where d is the cardinality of $\Xi(1)$. This follows form the fact that equivariant line bundles over X are parametrized by \mathbb{Z}^d (see [10]).

4. Kaneyama type description

In this section our main goal is to give a more combinatorial flavor to our classification. We fix an embedding of G in $GL(r,\mathbb{C})$ such that $K_0 := (\mathbb{C}^*)^r \cap G$ is a maximal torus of G. Inspired by the work of Kaneyama, we restrict to a smaller class of distinguished sections.

Definition 4.1. A distinguished section s of \mathcal{E}_{σ} is called a Kaneyama section if the image of the corresponding homomorphism ρ_s lies in K_0 .

Lemma 4.2. Every \mathcal{E}_{σ} has a Kaneyama section.

Proof. Let s be a distinguished section of \mathcal{E}_{σ} . The image of the corresponding homomorphism $\rho_s: T \to G$ is contained in some maximal torus K_{σ} of G. Since all maximal tori in G are conjugate, there exists $h_{\sigma} \in G$ be such that $h_{\sigma}K_{\sigma}h_{\sigma}^{-1} = K_0$. Then $h_{\sigma}\rho_s(t)h_{\sigma}^{-1} \in K_0$. Hence $s \cdot h_{\sigma}$ is a Kaneyama section.

For a Kaneyama section s_{σ} , the image $\rho_{\sigma}(t)$ is a diagonal matrix which we denote by

$$\xi^{\sigma}(t) = \operatorname{diag}(\xi_1^{\sigma}(t), \ldots, \xi_r^{\sigma}(t)).$$

Note that for different choices of Kaneyama sections s_{σ} , the images $\rho_{\sigma}(t)$ can vary up to conjugation (see Lemma 2.9), but they are all diagonal matrices.

Lemma 4.3. If the maximal torus K_0 is normal in G, then any T-equivariant holomorphic (or algebraic) principal G-bundle over a nonsingular toric variety admits an equivariant reduction of structure group to K_0 .

Proof. Let $\{s_{\sigma}\}$ be a collection of Kaneyama sections as σ varies over all maximal cones. Let $\xi^{\tau}(t)P(\tau,\sigma)\xi^{\sigma}(t)^{-1}$ be the corresponding transition maps. For each σ , we choose $g_{\sigma} \in G$ such that all $s_{\sigma} \cdot g_{\sigma}$'s pass through a common point in $\mathcal{E}(x_0)$. In other words, for these new sections, all $P(\tau,\sigma) = 1_G$. The new transition maps are $g_{\tau}^{-1}\xi^{\tau}(t)g_{\tau}g_{\sigma}^{-1}\xi^{\sigma}(t)^{-1}g_{\sigma}$. These are in K_0 as it is normal in G.

An analogue of the following lemma was proved in [7] in the context of vector bundles. The idea of our proof is the same as in [7], but we include it for clarity. The result is expected as the $T_{\sigma \cap \tau}$ -actions on the fibers of \mathcal{E}_{σ} and \mathcal{E}_{τ} at a fixed point $x_{\sigma \cap \tau} \in X_{\sigma} \cap X_{\tau}$ of $T_{\sigma \cap \tau}$ agree.

Lemma 4.4. Suppose σ and τ are cones of Ξ . Then there is a permutation γ of $\{1,\ldots,r\}$ such that $\eta(\xi_i^{\sigma}) = \eta(\xi_{\gamma(i)}^{\tau})$ for every $\eta \in \sigma \cap \tau \cap \Xi(1)$.

Proof. Recall that the transition function $\xi^{\tau}(t)P(\tau,\sigma)\xi^{\sigma}(t)^{-1}$ is regular on $X_{\sigma\cap\tau}$. Consider $P(\tau,\sigma)$ as a matrix and let $p_{ij}=P(\tau,\sigma)_{ij}$. Then $p_{ij}\xi_i^{\tau}(t)\xi_j^{\sigma}(t)^{-1}$ is regular on $X_{\sigma\cap\tau}$ for each (i,j).

Each ξ_i^{σ} may be considered as an element of the dual lattice M of N. Then regularity implies that

$$\eta(\xi_i^{\tau}) \ge \eta(\xi_j^{\sigma}) \tag{4.1}$$

for every $\eta \in \sigma \cap \tau \cap \Xi(1)$ whenever $p_{ij} \neq 0$.

Since

$$\det P(\tau, \sigma) = \sum_{\gamma \in S_r} \operatorname{sign}(\gamma) p_{1\gamma(1)} \dots p_{r\gamma(r)} \neq 0,$$

there exists a permutation γ such that $p_{i\gamma(i)} \neq 0$ for every i. Therefore,

$$\eta(\xi_i^{\tau}) \ge \eta(\xi_{\gamma(i)}^{\sigma})$$

for every i and every $\eta \in \sigma \cap \tau \cap \Xi(1)$.

Moreover since

$$\det\left[\xi^{\tau}(t)P(\tau,\sigma)\xi^{\sigma}(t)^{-1}\right] = \det P(\tau,\sigma)\prod_{i}\xi_{i}^{\tau}(t)\xi_{i}^{\sigma}(t)^{-1}$$

is a unit on $X_{\sigma \cap \tau}$, we have

$$\eta(\xi_1^{\tau} + \dots \xi_r^{\tau} - \xi_1^{\sigma} - \dots \xi_r^{\sigma}) = 0$$
(4.2)

for every $\eta \in \sigma \cap \tau \cap \Xi(1)$. Comparing (4.2) with (4.1), we have

$$\eta(\xi_i^\tau) = \eta(\xi_{\gamma(i)}^\sigma)$$

for every i and every η in $\sigma \cap \tau \cap \Xi(1)$.

Write

$$\xi^{\sigma} := (\xi_1^{\sigma}, \dots, \xi_r^{\sigma}) \in M^{\oplus r}.$$

Recall that $\rho_{\sigma}(t)$, and therefore $\xi^{\sigma}(t)$ must factor through the projection $\pi_{\sigma}: T \to T_{\sigma}$. There is a map $\pi_{\sigma}^*: M_{\sigma} \to M$, associated to π_{σ} , from the characters of T_{σ} to those of T. Then, ξ^{σ} lies in the image of $M_{\sigma}^{\oplus r}$ in $M^{\oplus r}$ under the map induced by π_{σ}^* . Note that this image is same as $M^{\oplus r}$ if σ is of top dimension, as $T_{\sigma} = T$ in this case.

Denote the set of maximal cones of Ξ by Ξ^* . Consider the following abstract data.

(1) A map

$$\xi: \Xi^* \to (\pi_\sigma^*(M_\sigma))^{\oplus r}$$

sending σ to $\xi^{\sigma} = (\xi_1^{\sigma}, \dots, \xi_r^{\sigma})$ such that for every pair of cones $\sigma, \tau \in \Xi(n)$, there exists a permutation γ such that

$$\eta(\xi_i^{\sigma}) = \eta(\xi_{\gamma(i)}^{\tau})$$

for every i and every $\eta \in \sigma \cap \tau \cap \Xi(1)$.

(2) A map

$$P:\Xi^*\times\Xi^*\to G\leq GL(r)$$

sending (τ, σ) to $P(\tau, \sigma)$ such that $P(\tau, \sigma)_{ij} \neq 0$ only if $\eta(\xi_i^{\tau}) \geq \eta(\xi_j^{\sigma})$ for every η in $(\tau \cap \sigma) \cap \Xi(1)$, and such that $P(\sigma, \sigma) = 1_G$ and

$$P(\tau, \sigma)P(\sigma, \delta)P(\tau, \delta) = 1_G$$

for every τ, σ, δ in Ξ^* .

(3) Write (ξ, P) for a collection $\{\xi^{\sigma}, P(\tau, \sigma)\}$ where σ, τ vary over Ξ^* . Two pairs (ξ, P) and (ξ', P') are said to be equivalent if there exists a permutation $\gamma \in S_r$, depending on σ , such that

$$(\xi_1^{\sigma},\ldots,\xi_r^{\sigma})=(\xi_{\gamma(1)}^{\prime\sigma},\ldots,\xi_{\gamma(r)}^{\prime\sigma})$$

for every $\sigma \in \Xi^*$, and if there exists

$$g:\Xi\to G$$

such that

$$P'(\tau, \sigma) = g(\tau)^{-1} P(\tau, \sigma) g(\sigma)$$

for every τ, σ in Ξ^* .

Theorem 4.5. Let X be a nonsingular toric variety defined by a fan Ξ of dimension n. The set of isomorphism classes of T-equivariant principal holomorphic (or algebraic) G-bundles on X is in bijective correspondence with the set of data (1) and (2) up to the equivalence (3).

Proof. Given a principal bundle, a choice of Kaneyama sections $\{s_{\sigma}\}$ associates a set of data (ξ, P) satisfying (1) and (2) to it. Note that $\{\xi_{i}^{\sigma} : 1 \leq i \leq r\}$ is the set of characters of the representation ρ_{σ} . Hence it is invariant under conjugation and therefore well-defined up to permutation under choice of different Kaneyama sections $s_{\sigma} \cdot g(\sigma)$. We have seen before that such a change would transform $\{P(\tau, \sigma)\}$ to $\{g(\tau)^{-1}P(\tau, \sigma)g(\sigma)\}$. Same arguments as in the proof of Theorem 3.2 imply that

this association sends an isomorphism class of bundles to an isomorphism class of data.

On the other hand, given a set of data (ξ, P) satisfying (1) and (2), we regard ξ^{σ} as ρ_{σ} and $P(\tau, \sigma)$ as transition map at a point $x_0 \in O$ to construct a bundle $E(\{\rho, P\})$. The rest of the proof proceeds as in Theorem 3.2.

Given ξ , for every $\sigma \in \Xi(n)$, we have a map

$$m_{\sigma}: \sigma \bigcap \Xi(1) \to \mathbb{Z}^r$$

defined by $m_{\sigma}(\eta)_i = \eta(\xi_i^{\sigma})$. Note that as σ is smooth one may recover ξ^{σ} from m_{σ} . A strategy of Kaneyama is to reformulate the equivalence (3) in terms of m_{σ} . It turns out as follows.

(3') Two pairs (ξ, P) and (ξ', P') are said to be equivalent if there exists a permutation $\gamma = \gamma(\sigma) \in S(r)$ such that

$$(m_{\sigma}(\eta)_1,\ldots,m_{\sigma}(\eta)_r)=(m'_{\sigma}(\eta)_{\gamma(1)},\ldots,m'_{\sigma}(\eta)_{\gamma(r)})$$

for every $\sigma \in \Xi(n)$ and $\eta \in \sigma \cap \Xi(1)$, and if there exists

$$g:\Xi(n)\to G$$

such that

$$P'(\tau, \sigma) = g(\tau)^{-1} P(\tau, \sigma) g(\sigma)$$

for every τ , σ in $\Xi(n)$.

The proof of the following result is straight-forward.

Theorem 4.6. Let X be a nonsingular toric variety defined by a fan Ξ of dimension n, such that every maximal cone of Ξ has dimension n. Then, the set of isomorphism classes of T-equivariant holomorphic (or algebraic) principal G-bundles on X is in bijective correspondence with the set of data (1) and (2) up to the equivalence (3').

Remark 4.7. A conjecture of Hartshorne [4] says that all vector bundles of rank two over \mathbb{P}^n should split if $n \geq 7$. This is still open. In [8], the following equivariant analogue is discussed: Any T-equivariant vector bundle of rank r on \mathbb{P}^n splits if r < n. An attempt to reproduce the arguments given there runs into the problem that conjugation by an elementary matrix does not preserve the diagonal matrices. A stronger result has recently been proved in an article of Ilten and Süss [6] using Klyachko-type filtrations [9]. It would be interesting to investigate if their result can be generalized to equivariant principal bundles.

Acknowledgement. It is a pleasure to thank V. Balaji for telling us about the work of Heinzner and Kutzschebauch, and Pralay Chatterjee for many useful discussions. We thank an anonymous referee for his/her valuable comments on an earlier draft. The first-named author is supported by a J. C. Bose fellowship. The last-named author is supported by a FAPA grant from the Universidad de los Andes.

REFERENCES

- [1] A. Dey and M. Poddar: Equivariant Abelian principal bundles on nonsingular toric varieties, *Bull. Sci. Math.* (to appear).
- [2] W. Fulton: Introduction to Toric Varieties, Annals of Mathematics Studies, 131, Princeton University Press, Princeton, 1993.
- [3] I. Hambleton, Ian and J.-C. Hausmann: Equivariant bundles and isotropy representations, *Groups Geom. Dyn.* 4 (2010), no. 1, 127-162.
- [4] R. Hartshorne: Varieties of small codimension in projective space, Bull. Amer. Math. Soc. 80 (1974), 1017–1032.
- [5] P. Heinzner and F. Kutzschebauch: An equivariant version of Grauert's Oka principle, *Invent. Math.* **119** (1995), 317–346.
- [6] N. Ilten and H. Süss: Equivariant vector bundles on T-varieties, arXiv:1305.0024 Transform. Groups (to appear).
- [7] T. Kaneyama: On equivariant vector bundles on an almost homogeneous variety, *Nagoya Math. Jour.* **57** (1975), 65–86.
- [8] T. Kaneyama: Torus-equivariant vector bundles on projective spaces, *Nagoya Math. Jour.* **111** (1988), 25–40.
- [9] A. A. Klyachko: Equivariant bundles over toric varieties, Izv. Akad. Nauk SSSR Ser. Mat. 53 (1989), 1001–1039, 1135; translation in Math. USSR-Izv. 35 (1990), 337–375
- [10] T. Oda: Convex bodies and algebraic geometry. An introduction to the theory of toric varieties, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 15. Springer-Verlag, Berlin, 1988.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, MUMBAI, INDIA E-mail address: indranil@math.tifr.res.in

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY-MADRAS, CHENNAI, INDIA

E-mail address: arijitdey@gmail.com

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE LOS ANDES, BOGOTÁ, COLOMBIA; AND, DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, NORTHERN CYPRUS CAMPUS, KALKANLI, GUZELYURT, KKTC, MERSIN 10 TURKEY

E-mail address: mainakp@gmail.com