

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 3-({[(1-Phenylethyl)sulfanyl]methane-thioyl}sulfanyl)propanoic acid

M. Kannan, V. Ramkumar and R. Dhamodharan\*

Department of Chemistry, IIT Madras, Chennai, Tamil Nadu, India  
Correspondence e-mail: damo@iitm.ac.in


Received 27 October 2011; accepted 7 November 2011

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.036;  $wR$  factor = 0.095; data-to-parameter ratio = 19.0.

In the title compound,  $\text{C}_{12}\text{H}_{14}\text{O}_2\text{S}_3$ , a chain transfer agent (CTA) used in polymerization, the dihedral angle between the aromatic ring and the  $\text{CS}_3$  grouping is  $84.20(10)^\circ$ . In the crystal, carboxylic acid inversion dimers linked by pairs of  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonds generate  $R_2^2(8)$  loops.

## Related literature

For background to chain transfer agents, see: Chong *et al.* (1999); Coady *et al.* (2008). For a related structure, see: Kannan *et al.* (2010).



## Experimental

## Crystal data

$\text{C}_{12}\text{H}_{14}\text{O}_2\text{S}_3$   
 $M_r = 286.41$   
 Monoclinic,  $P2_1/c$   
 $a = 13.6280(8)$  Å  
 $b = 10.2908(5)$  Å

$c = 10.7299(5)$  Å  
 $\beta = 113.039(2)^\circ$   
 $V = 1384.77(12)$  Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation

$\mu = 0.52$  mm<sup>-1</sup>  
 $T = 298$  K

 $0.42 \times 0.28 \times 0.22$  mm

## Data collection

Bruker APEXII CCD area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2004)  
 $T_{\min} = 0.811$ ,  $T_{\max} = 0.894$

9109 measured reflections  
 3020 independent reflections  
 2224 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.017$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$   
 $wR(F^2) = 0.095$   
 $S = 1.03$   
 3020 reflections  
 159 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.32$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.27$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O2}-\text{H1O}\cdots\text{O1}^i$	0.81 (3)	1.85 (3)	2.651 (2)	177 (3)

Symmetry code: (i)  $-x, -y + 3, -z$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2004); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97.

The authors acknowledge the Department of Chemistry, IIT Madras, for the data collection. MK thanks the CSIR, India, for a fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6480).

## References

- Bruker (2004). APEX2, SAINT-Plus and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Chong, Y. K., Le, T. P. T., Moad, G., Rizzardo, E. & Thang, S. H. (1999). *Macromolecules*, **32**, 2071–2074.  
 Coady, D. J., Norris, B. C., Lynch, V. M. & Bielawski, C. W. (2008). *Macromolecules*, **41**, 3775–3778.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Kannan, M., Ramkumar, V. & Dhamodharan, R. (2010). *Acta Cryst.* **E66**, o1382.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

## supporting information

*Acta Cryst.* (2011). E67, o3352 [https://doi.org/10.1107/S1600536811046873]

### 3-([(1-Phenylethyl)sulfanyl]methanethioyl)sulfanyl)propanoic acid

M. Kannan, V. Ramkumar and R. Dhamodharan

#### S1. Comment

The title compound  $C_{12}H_{14}S_3O_2$  is a carbanotrithioate. It can be used as a chain transfer agent (CTA) in RAFT polymerization (Chong *et al.*, 1999) to control the polymerization and it will produce carbanotrithionate end-terminated polymers. Very few single-crystal XRD data are available for CTAs, because most of them are liquids (Coady *et al.*, 2008). Recently, we have reported the single-crystal data of a multi-functional CTA, which can be used for the synthesis of star polymers. Carbanotrithioate CTA is suitable for the polymerization of styrene, acrylates and methacrylates. With appropriate choice of the CTA (RAFT agent) and reaction conditions, RAFT polymerization can be successfully used to produce polymers of narrow polydispersity with predetermined molecular weights. Moreover, the polymers obtained by the RAFT process can be chain extended or used as precursors to synthesize stimuli responsive block copolymers by the addition of further monomer(s). The title compound will result in carboxylic acid end-terminated polymer; this functionality can be further modified and utilized for making block copolymers by reacting it with another homopolymer.

The compound  $C_{12}H_{14}S_3O_2$  is stabilized by a O—H $\cdots$ O interaction with  $R_2^2(8)$  graph set motif.

#### S2. Experimental

The title compound, was prepared by adding 3-mercapto propanoic acid (1.00 g, 7.35 mmol) to a stirred suspension of  $K_3PO_4$  (1.72 g, 8.09 mmol) in acetone (20 ml) over a period of ten minutes.  $CS_2$  (1.68 g, 22.06 mmol) was added upon which the solution turned bright yellow. After stirring for ten minutes 1-bromo ethyl benzene (1.26 g, 7.35 mmol) was added and an instant precipitation of KBr was noted. After stirring for three hours the suspension was filtered and the cake was rinsed with acetone ( $2 \times 20$  ml). After removing the solvent from the filtrate under reduced pressure the resulting yellow residue was purified by column chromatography on silica using a petroleum ether/ethyl acetate gradient to yield light yellow solid (96%) that crystallized to form light yellow blocks.

#### S3. Refinement

All hydrogen atoms were fixed geometrically and allowed to ride on the parent carbon atoms, with aromatic C—H = 0.93 Å, methyl C—H = 0.96 Å and methylene C—H = 0.97 Å. The displacement parameters were set for phenyl and methylene H atoms at  $U_{iso}(H) = 1.2U_{eq}(C)$  and methyl H atoms at  $U_{iso}(H) = 1.5U_{eq}(C)$ .




Figure 1

The molecular structure of the title molecule with atoms represented as 30% probability ellipsoids.

### 3-((1-Phenylethyl)sulfanyl)propanoic acid

#### Crystal data

$C_{12}H_{14}O_2S_3$

$M_r = 286.41$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2_1/c$

$a = 13.6280$  (8) Å

$b = 10.2908$  (5) Å

$c = 10.7299$  (5) Å

$\beta = 113.039$  (2)°

$V = 1384.77$  (12) Å<sup>3</sup>

$Z = 4$

$F(000) = 600$

$D_x = 1.374$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 4284 reflections

$\theta = 2.6$ – $27.2$ °

$\mu = 0.52$  mm<sup>-1</sup>

$T = 298$  K

Block, light yellow

$0.42 \times 0.28 \times 0.22$  mm

#### Data collection

Bruker APEXII CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2004)

$T_{\min} = 0.811$ ,  $T_{\max} = 0.894$

9109 measured reflections

3020 independent reflections

2224 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 28.5$ °,  $\theta_{\min} = 2.6$ °

$h = -17 \rightarrow 18$

$k = -12 \rightarrow 12$

$l = -14 \rightarrow 12$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.095$

$S = 1.03$

3020 reflections

159 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H atoms treated by a mixture of independent  
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0372P)^2 + 0.5628P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.32$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.27$  e Å<sup>-3</sup>

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.34830 (19)	0.5248 (2)	0.1937 (3)	0.0664 (6)
H1	0.3225	0.5785	0.1180	0.080*
C2	0.4227 (2)	0.4306 (3)	0.2022 (4)	0.0874 (9)
H2	0.4469	0.4216	0.1328	0.105*
C3	0.4606 (2)	0.3509 (3)	0.3116 (4)	0.0948 (11)
H3	0.5103	0.2870	0.3168	0.114*
C4	0.4260 (2)	0.3645 (3)	0.4129 (4)	0.0912 (10)
H4	0.4519	0.3095	0.4875	0.109*
C5	0.3523 (2)	0.4598 (2)	0.4070 (3)	0.0684 (6)
H5	0.3302	0.4692	0.4783	0.082*
C6	0.31156 (16)	0.54066 (18)	0.2959 (2)	0.0492 (5)
C7	0.23156 (16)	0.64611 (17)	0.2845 (2)	0.0477 (5)
H7	0.1806	0.6504	0.1901	0.057*
C8	0.1697 (2)	0.6301 (3)	0.3737 (3)	0.0832 (8)
H8A	0.1330	0.5482	0.3546	0.125*
H8B	0.1187	0.6993	0.3559	0.125*
H8C	0.2180	0.6327	0.4671	0.125*
C9	0.21603 (15)	0.91964 (17)	0.24871 (19)	0.0421 (4)
C10	0.18484 (18)	1.18835 (19)	0.2244 (2)	0.0553 (5)
H10A	0.1186	1.1527	0.2231	0.066*
H10B	0.2046	1.2608	0.2872	0.066*
C11	0.16642 (16)	1.23796 (18)	0.0855 (2)	0.0479 (5)
H11A	0.2342	1.2624	0.0825	0.057*
H11B	0.1359	1.1692	0.0199	0.057*
C12	0.09317 (15)	1.35291 (18)	0.0482 (2)	0.0444 (4)
O1	0.04244 (14)	1.38506 (15)	0.11389 (17)	0.0706 (5)
O2	0.08929 (15)	1.41368 (17)	-0.05809 (18)	0.0708 (5)
S1	0.30835 (4)	0.79721 (5)	0.32472 (6)	0.05737 (18)
S2	0.09018 (4)	0.90297 (6)	0.15929 (6)	0.06006 (18)
S3	0.28670 (5)	1.06576 (5)	0.28523 (7)	0.06266 (19)
H1O	0.051 (3)	1.477 (3)	-0.073 (3)	0.111 (12)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0664 (15)	0.0545 (13)	0.0817 (17)	0.0057 (11)	0.0326 (13)	-0.0060 (12)
C2	0.0686 (18)	0.0723 (18)	0.125 (3)	0.0037 (14)	0.0424 (18)	-0.0282 (18)
C3	0.0550 (17)	0.0490 (15)	0.163 (3)	0.0051 (12)	0.023 (2)	-0.0112 (18)
C4	0.0649 (18)	0.0499 (15)	0.127 (3)	-0.0007 (13)	0.0035 (18)	0.0247 (16)
C5	0.0650 (15)	0.0477 (13)	0.0813 (17)	-0.0060 (11)	0.0166 (13)	0.0118 (12)
C6	0.0460 (12)	0.0316 (10)	0.0666 (14)	-0.0063 (8)	0.0182 (10)	-0.0023 (9)
C7	0.0498 (12)	0.0357 (10)	0.0610 (13)	-0.0039 (8)	0.0256 (10)	0.0005 (9)
C8	0.098 (2)	0.0674 (16)	0.115 (2)	0.0002 (15)	0.0750 (19)	0.0082 (15)
C9	0.0449 (11)	0.0387 (10)	0.0425 (11)	0.0042 (8)	0.0170 (9)	0.0011 (8)
C10	0.0641 (14)	0.0361 (10)	0.0645 (14)	0.0124 (9)	0.0240 (11)	0.0065 (9)
C11	0.0471 (11)	0.0399 (10)	0.0585 (13)	0.0088 (8)	0.0228 (10)	0.0041 (9)
C12	0.0457 (11)	0.0380 (10)	0.0515 (12)	0.0057 (8)	0.0214 (10)	0.0048 (8)
O1	0.0899 (12)	0.0649 (10)	0.0778 (11)	0.0397 (9)	0.0553 (10)	0.0286 (8)
O2	0.0879 (13)	0.0675 (11)	0.0748 (12)	0.0384 (10)	0.0510 (10)	0.0314 (9)
S1	0.0441 (3)	0.0327 (3)	0.0829 (4)	0.0021 (2)	0.0114 (3)	0.0055 (2)
S2	0.0411 (3)	0.0620 (3)	0.0680 (4)	0.0046 (2)	0.0115 (3)	-0.0083 (3)
S3	0.0507 (3)	0.0354 (3)	0.0857 (5)	0.0032 (2)	0.0092 (3)	0.0112 (3)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C1—C2	1.380 (3)	C8—H8B	0.9600
C1—C6	1.383 (3)	C8—H8C	0.9600
C1—H1	0.9300	C9—S2	1.614 (2)
C2—C3	1.357 (5)	C9—S1	1.7415 (19)
C2—H2	0.9300	C9—S3	1.7455 (19)
C3—C4	1.351 (5)	C10—C11	1.500 (3)
C3—H3	0.9300	C10—S3	1.799 (2)
C4—C5	1.388 (4)	C10—H10A	0.9700
C4—H4	0.9300	C10—H10B	0.9700
C5—C6	1.380 (3)	C11—C12	1.498 (2)
C5—H5	0.9300	C11—H11A	0.9700
C6—C7	1.509 (3)	C11—H11B	0.9700
C7—C8	1.512 (3)	C12—O1	1.211 (2)
C7—S1	1.8291 (19)	C12—O2	1.283 (2)
C7—H7	0.9800	O2—H1O	0.81 (3)
C8—H8A	0.9600		
C2—C1—C6	121.0 (3)	H8A—C8—H8B	109.5
C2—C1—H1	119.5	C7—C8—H8C	109.5
C6—C1—H1	119.5	H8A—C8—H8C	109.5
C3—C2—C1	120.2 (3)	H8B—C8—H8C	109.5
C3—C2—H2	119.9	S2—C9—S1	127.34 (11)
C1—C2—H2	119.9	S2—C9—S3	126.14 (11)
C4—C3—C2	119.9 (3)	S1—C9—S3	106.51 (11)
C4—C3—H3	120.0	C11—C10—S3	113.87 (14)

C2—C3—H3	120.0	C11—C10—H10A	108.8
C3—C4—C5	120.8 (3)	S3—C10—H10A	108.8
C3—C4—H4	119.6	C11—C10—H10B	108.8
C5—C4—H4	119.6	S3—C10—H10B	108.8
C6—C5—C4	120.3 (3)	H10A—C10—H10B	107.7
C6—C5—H5	119.9	C12—C11—C10	111.69 (16)
C4—C5—H5	119.9	C12—C11—H11A	109.3
C5—C6—C1	117.8 (2)	C10—C11—H11A	109.3
C5—C6—C7	122.5 (2)	C12—C11—H11B	109.3
C1—C6—C7	119.66 (19)	C10—C11—H11B	109.3
C6—C7—C8	115.77 (18)	H11A—C11—H11B	107.9
C6—C7—S1	105.29 (13)	O1—C12—O2	123.36 (18)
C8—C7—S1	110.65 (16)	O1—C12—C11	122.18 (17)
C6—C7—H7	108.3	O2—C12—C11	114.46 (16)
C8—C7—H7	108.3	C12—O2—H1O	112 (2)
S1—C7—H7	108.3	C9—S1—C7	105.23 (9)
C7—C8—H8A	109.5	C9—S3—C10	104.07 (10)
C7—C8—H8B	109.5		
C6—C1—C2—C3	0.4 (4)	C1—C6—C7—S1	-76.6 (2)
C1—C2—C3—C4	-0.5 (4)	S3—C10—C11—C12	-171.50 (14)
C2—C3—C4—C5	-0.3 (4)	C10—C11—C12—O1	-11.6 (3)
C3—C4—C5—C6	1.2 (4)	C10—C11—C12—O2	168.19 (19)
C4—C5—C6—C1	-1.3 (3)	S2—C9—S1—C7	-1.18 (16)
C4—C5—C6—C7	-179.8 (2)	S3—C9—S1—C7	179.94 (9)
C2—C1—C6—C5	0.5 (3)	C6—C7—S1—C9	156.85 (14)
C2—C1—C6—C7	179.0 (2)	C8—C7—S1—C9	-77.38 (19)
C5—C6—C7—C8	-20.6 (3)	S2—C9—S3—C10	8.69 (16)
C1—C6—C7—C8	160.9 (2)	S1—C9—S3—C10	-172.41 (10)
C5—C6—C7—S1	101.9 (2)	C11—C10—S3—C9	-97.04 (17)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H1O···O1 <sup>i</sup>	0.81 (3)	1.85 (3)	2.651 (2)	177 (3)

Symmetry code: (i)  $-x, -y+3, -z$ .