

# Crystal structure of aquadioxido-(2-[(2-oxidoethyl)imino]methyl]phenolato- $\kappa^3 O, N, O'$ )molybdenum(VI)

Sowmianarayanan Parimala<sup>a</sup> and Parasuraman Selvam<sup>a,b,c\*</sup>

<sup>a</sup>National Centre for Catalysis Research and Department of Chemistry, Indian Institute of Technology-Madras, Chennai 600 036, India, <sup>b</sup>New Industry Creation Hatchery Center, Tohoku University, Sendai 980 8579, Japan, and <sup>c</sup>School of Science and Health, University of Western Australia, Sydney, Penrith, NSW 275, Australia.

\*Correspondence e-mail: selvam@itm.ac.in

Received 9 January 2015; accepted 20 January 2015

Edited by M. Weil, Vienna University of Technology, Austria


The mononuclear title complex,  $[\text{Mo}(\text{C}_9\text{H}_9\text{NO}_2)\text{O}_2(\text{H}_2\text{O})]$ , contains an Mo(VI) atom in a distorted octahedral coordination sphere defined by an Mo=O and an Mo—(OH<sub>2</sub>) bond to the axial ligands and two Mo—O bonds to phenolate and alcoholate O atoms, another Mo=O bond and one Mo—N bond to the imino N atom in the equatorial plane. The five-membered metalla-ring shows an envelope conformation. In the crystal, individual molecules are connected into a layered arrangement parallel to (100) by means of O—H...O hydrogen bonds involving the water molecule as a donor group and the O atoms of neighbouring complexes as acceptor atoms. These interactions lead to the formation of a three-dimensional network.

**Keywords:** crystal structure; dioxidomolybdenum(VI) complex; hydrogen bonding.

**CCDC reference:** 1044382

## 1. Related literature

For dioxidomolybdenum complexes used as potential oxidation catalysts for the epoxidation of alkenes, see: Sakthivel *et al.* (2005); Masteri-Farahani *et al.* (2006). For chiral molybdenum complexes, see: Burke (2008); Kühn *et al.* (2005). These compounds are good catalysts for the oxidation of organic compounds, see: Rayati *et al.* (2012). For heterogenization of polymer-supported molybdenum complexes, see: Sherrington *et al.* (2000); Maurya (2012), and for molybdenum systems on silica supports, see: Tangestaninejad *et al.* (2008).



## 2. Experimental

### 2.1. Crystal data

$[\text{Mo}(\text{C}_9\text{H}_9\text{NO}_2)\text{O}_2(\text{H}_2\text{O})]$   
 $M_r = 309.13$   
 Monoclinic,  $P2_1/c$   
 $a = 14.9710(3) \text{ \AA}$   
 $b = 6.7026(1) \text{ \AA}$   
 $c = 10.8673(2) \text{ \AA}$   
 $\beta = 99.486(1)^\circ$

$V = 1075.56(3) \text{ \AA}^3$   
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.22 \text{ mm}^{-1}$   
 $T = 296 \text{ K}$   
 $0.25 \times 0.16 \times 0.10 \text{ mm}$

### 2.2. Data collection

Bruker APEXII CCD  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2012)  
 $T_{\min} = 0.813$ ,  $T_{\max} = 0.934$

8837 measured reflections  
 2392 independent reflections  
 2263 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.012$

### 2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.016$   
 $wR(F^2) = 0.042$   
 $S = 1.10$   
 2392 reflections  
 153 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.31 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.30 \text{ e \AA}^{-3}$

**Table 1**

Selected bond lengths ( $\text{\AA}$ ).

Mo1—O5	1.6902 (14)	Mo1—O2	1.9446 (12)
Mo1—O4	1.7160 (13)	Mo1—N1	2.2652 (14)
Mo1—O1	1.9438 (12)	Mo1—O3	2.3259 (14)

**Table 2**

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D\text{—}H\cdots A$	$D\text{—}H$	$H\cdots A$	$D\cdots A$	$D\text{—}H\cdots A$
O3—H1O...O1 <sup>i</sup>	0.73 (2)	1.97 (3)	2.6656 (19)	161 (3)
O3—H2O...O4 <sup>ii</sup>	0.78 (3)	2.07 (3)	2.8425 (19)	173 (3)

Symmetry codes: (i)  $-x + 1, -y, -z + 1$ ; (ii)  $x, -y + \frac{1}{2}, z - \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2012); cell refinement: *SAINTE* (Bruker, 2012); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2015); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *publCIF* (Westrip, 2010).

### Acknowledgements

The authors thank the Department of Science and Technology (DST), Government of India, for funding the National Centre for Catalysis Research (NCCR), IIT-Madras. They also thank Mr V. Ramkumar and Dr R. Jagan for the data collection and technical assistance in the preparation of the manuscript.

---

Supporting information for this paper is available from the IUCr electronic archives (Reference: WM5114).

---

### References

- Bruker (2012). *APEX2*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burke, A. (2008). *Coord. Chem. Rev.* **252**, 170–175.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Kühn, F. E. J., Zhao, J. & Herrmann, W. A. (2005). *Tetrahedron Asymmetry*, **16**, 3469–3479.
- Masteri-Farahani, M., Farzaneh, F. & Ghandi, M. J. (2006). *J. Mol. Catal. A Chem.* **248**, 53–60.
- Maurya, R. (2012). *Curr. Org. Chem.* **16**, 73–88.
- Rayati, S., Rafiee, N. & Wojtczak, A. (2012). *Inorg. Chim. Acta*, **386**, 27–35.
- Sakthivel, A., Zhao, J., Raudaschl-Sieber, G., Hanzlik, M., Chiang, A. S. T. & Kühn, F. E. (2005). *Appl. Catal. Gen.* **281**, 267–273.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sheldrick, G. M. (2015). *Acta Cryst.* **C71**, 3–8.
- Sherrington, D. C. (2000). *Catal. Today*, **57**, 87–104.
- Tangestaninejad, S., Moghadam, M., Mirkhani, V., Mohammadpoor-Baltork, I. & Ghani, K. (2008). *J. Iran Chem. Soc.* **5**, s71–S79.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## supporting information

*Acta Cryst.* (2015). E71, m35–m36 [doi:10.1107/S2056989015001231]

## Crystal structure of aquadioxido(2-[[2-(2-oxidoethyl)imino]methyl]phenolato- $\kappa^3O,N,O'$ )molybdenum(VI)


Sowmianarayanan Parimala and Parasuraman Selvam

### S1. Experimental

Molybdenyl acetylacetonate ( $\text{MoO}_2(\text{acac})_2$ ) (4.03 g, 0.012 mol) dissolved in methanol (20 ml) was added to a refluxing solution of salicylaldehyde (2.62 ml, 0.012 mol) and ethanolamine (1.5 ml, 0.012 mol) in ethanol (30 ml). The mixture was refluxed for five hours, and the solvent removed under vacuum at room temperature. The resulting yellow solution was filtered, evaporated slowly, to yield yellow crystals. The crystals were purified by washing with ethanol/methanol mixture and dried at room temperature. The obtained crystals have incorporated water. The used solvents ethanol and methanol have not been dried prior to the reaction and thus contain water. Another source of water is the condensation reaction between salicylaldehyde and ethanolamine.


### S2. Refinement

All H atoms were identified from difference electron density maps. However, C-bound H atoms were treated as riding with C—H = 0.97 Å for ( $\text{CH}_2$ ) and C—H = 0.93 Å for aromatic H atoms, both with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}$ . The H atoms of the water molecule were refined freely.



**Figure 1**

The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level.

**Figure 2**

Unit-cell packing diagram of the title compound with hydrogen bonds shown as dashed lines. Hydrogen atoms not involved in hydrogen bonding are omitted for clarity.

### Aquadioxido(2-[(2-oxidoethyl)imino]methyl)phenolato- $\kappa^3O,N,O'$ molybdenum(VI)

#### Crystal data

[Mo(C<sub>9</sub>H<sub>9</sub>NO<sub>2</sub>)O<sub>2</sub>(H<sub>2</sub>O)]

$M_r = 309.13$

Monoclinic,  $P2_1/c$

$a = 14.9710(3) \text{ \AA}$

$b = 6.7026(1) \text{ \AA}$

$c = 10.8673(2) \text{ \AA}$

$\beta = 99.486(1)^\circ$

$V = 1075.56(3) \text{ \AA}^3$

$Z = 4$

$F(000) = 616$

$D_x = 1.909 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6907 reflections

$\theta = 2.8\text{--}27.2^\circ$

$\mu = 1.22 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Block, yellow

$0.25 \times 0.16 \times 0.10 \text{ mm}$

#### Data collection

Bruker APEXII CCD

diffractometer

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(SADABS; Bruker, 2012)

$T_{\min} = 0.813$ ,  $T_{\max} = 0.934$

8837 measured reflections

2392 independent reflections

2263 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$

$\theta_{\max} = 27.2^\circ$ ,  $\theta_{\min} = 1.4^\circ$   
 $h = -19 \rightarrow 18$

$k = -8 \rightarrow 8$   
 $l = -13 \rightarrow 13$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.016$   
 $wR(F^2) = 0.042$   
 $S = 1.10$   
 2392 reflections  
 153 parameters  
 0 restraints

Hydrogen site location: mixed  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0163P)^2 + 0.8254P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.31 \text{ e } \text{Å}^{-3}$   
 $\Delta\rho_{\min} = -0.30 \text{ e } \text{Å}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{Å}^2$ )*

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.37800 (15)	-0.3126 (3)	0.4461 (2)	0.0394 (5)
H1A	0.3281	-0.3758	0.4778	0.047*
H1B	0.4281	-0.4060	0.4541	0.047*
C2	0.34936 (14)	-0.2553 (3)	0.31132 (19)	0.0384 (5)
H2A	0.4018	-0.2223	0.2734	0.046*
H2B	0.3169	-0.3641	0.2649	0.046*
C3	0.22629 (12)	-0.0467 (3)	0.22157 (16)	0.0288 (4)
H3	0.2174	-0.1379	0.1562	0.035*
C4	0.16677 (11)	0.1238 (3)	0.21392 (16)	0.0267 (4)
C5	0.09354 (13)	0.1343 (3)	0.11473 (18)	0.0371 (4)
H5	0.0846	0.0313	0.0567	0.044*
C6	0.03507 (13)	0.2930 (4)	0.10178 (19)	0.0427 (5)
H6	-0.0136	0.2961	0.0365	0.051*
C7	0.04887 (13)	0.4486 (4)	0.1865 (2)	0.0404 (5)
H7	0.0090	0.5561	0.1780	0.049*
C8	0.12118 (13)	0.4458 (3)	0.28334 (18)	0.0331 (4)
H8	0.1304	0.5525	0.3387	0.040*
C9	0.18051 (11)	0.2837 (3)	0.29870 (15)	0.0244 (3)
Mo1	0.32157 (2)	0.08935 (2)	0.49415 (2)	0.02134 (5)
N1	0.29025 (10)	-0.0807 (2)	0.31202 (14)	0.0254 (3)
O1	0.40476 (8)	-0.13539 (19)	0.51476 (12)	0.0286 (3)
O2	0.25056 (8)	0.29088 (18)	0.39282 (11)	0.0275 (3)
O3	0.42590 (10)	0.2153 (2)	0.37755 (13)	0.0306 (3)
O4	0.37757 (10)	0.2275 (2)	0.61586 (12)	0.0372 (3)
O5	0.23417 (9)	-0.0248 (2)	0.54594 (13)	0.0392 (3)
H1O	0.4715 (17)	0.176 (4)	0.395 (2)	0.037 (7)*
H2O	0.4144 (17)	0.221 (4)	0.305 (3)	0.048 (7)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0454 (11)	0.0219 (9)	0.0463 (12)	0.0080 (8)	-0.0058 (9)	-0.0057 (8)
C2	0.0417 (11)	0.0331 (10)	0.0382 (11)	0.0118 (8)	-0.0003 (8)	-0.0154 (8)
C3	0.0325 (9)	0.0312 (9)	0.0219 (8)	-0.0039 (7)	0.0024 (7)	-0.0059 (7)
C4	0.0239 (8)	0.0331 (9)	0.0223 (8)	-0.0025 (7)	0.0019 (6)	0.0033 (7)
C5	0.0344 (10)	0.0460 (12)	0.0274 (9)	-0.0061 (9)	-0.0045 (8)	0.0021 (9)
C6	0.0301 (10)	0.0601 (14)	0.0341 (10)	-0.0005 (9)	-0.0064 (8)	0.0154 (10)
C7	0.0325 (10)	0.0508 (13)	0.0384 (11)	0.0142 (9)	0.0068 (8)	0.0183 (10)
C8	0.0351 (10)	0.0363 (10)	0.0286 (9)	0.0097 (8)	0.0075 (7)	0.0061 (8)
C9	0.0231 (8)	0.0295 (9)	0.0207 (8)	0.0010 (7)	0.0038 (6)	0.0059 (7)
Mo1	0.02586 (8)	0.02184 (8)	0.01569 (8)	0.00375 (5)	0.00161 (5)	-0.00128 (5)
N1	0.0283 (7)	0.0234 (7)	0.0241 (7)	0.0011 (6)	0.0031 (6)	-0.0047 (6)
O1	0.0297 (6)	0.0241 (6)	0.0295 (6)	0.0054 (5)	-0.0023 (5)	-0.0007 (5)
O2	0.0306 (6)	0.0243 (6)	0.0247 (6)	0.0054 (5)	-0.0035 (5)	-0.0023 (5)
O3	0.0250 (7)	0.0404 (8)	0.0251 (7)	0.0017 (6)	0.0007 (5)	0.0052 (6)
O4	0.0479 (8)	0.0381 (8)	0.0219 (6)	0.0062 (6)	-0.0049 (6)	-0.0085 (6)
O5	0.0370 (7)	0.0451 (8)	0.0382 (8)	0.0037 (6)	0.0144 (6)	0.0072 (7)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

C1—O1	1.425 (2)	C6—H6	0.9300
C1—C2	1.507 (3)	C7—C8	1.380 (3)
C1—H1A	0.9700	C7—H7	0.9300
C1—H1B	0.9700	C8—C9	1.396 (2)
C2—N1	1.468 (2)	C8—H8	0.9300
C2—H2A	0.9700	C9—O2	1.3397 (19)
C2—H2B	0.9700	Mo1—O5	1.6902 (14)
C3—N1	1.275 (2)	Mo1—O4	1.7160 (13)
C3—C4	1.443 (3)	Mo1—O1	1.9438 (12)
C3—H3	0.9300	Mo1—O2	1.9446 (12)
C4—C9	1.406 (3)	Mo1—N1	2.2652 (14)
C4—C5	1.407 (2)	Mo1—O3	2.3259 (14)
C5—C6	1.370 (3)	O3—H1O	0.73 (2)
C5—H5	0.9300	O3—H2O	0.78 (3)
C6—C7	1.384 (3)		
O1—C1—C2	107.86 (16)	C7—C8—H8	119.8
O1—C1—H1A	110.1	C9—C8—H8	119.8
C2—C1—H1A	110.1	O2—C9—C8	117.79 (16)
O1—C1—H1B	110.1	O2—C9—C4	122.62 (15)
C2—C1—H1B	110.1	C8—C9—C4	119.56 (16)
H1A—C1—H1B	108.4	O5—Mo1—O4	107.11 (7)
N1—C2—C1	105.88 (15)	O5—Mo1—O1	97.32 (6)
N1—C2—H2A	110.6	O4—Mo1—O1	96.15 (6)
C1—C2—H2A	110.6	O5—Mo1—O2	97.01 (6)
N1—C2—H2B	110.6	O4—Mo1—O2	102.33 (6)

C1—C2—H2B	110.6	O1—Mo1—O2	152.09 (5)
H2A—C2—H2B	108.7	O5—Mo1—N1	90.18 (6)
N1—C3—C4	124.35 (16)	O4—Mo1—N1	161.73 (6)
N1—C3—H3	117.8	O1—Mo1—N1	75.35 (5)
C4—C3—H3	117.8	O2—Mo1—N1	80.78 (5)
C9—C4—C5	118.37 (17)	O5—Mo1—O3	166.38 (6)
C9—C4—C3	122.93 (15)	O4—Mo1—O3	86.41 (6)
C5—C4—C3	118.66 (17)	O1—Mo1—O3	82.47 (5)
C6—C5—C4	121.4 (2)	O2—Mo1—O3	78.05 (5)
C6—C5—H5	119.3	N1—Mo1—O3	76.56 (5)
C4—C5—H5	119.3	C3—N1—C2	121.04 (15)
C5—C6—C7	119.60 (18)	C3—N1—Mo1	127.21 (12)
C5—C6—H6	120.2	C2—N1—Mo1	111.56 (11)
C7—C6—H6	120.2	C1—O1—Mo1	117.80 (11)
C8—C7—C6	120.62 (19)	C9—O2—Mo1	133.92 (11)
C8—C7—H7	119.7	Mo1—O3—H1O	114.5 (19)
C6—C7—H7	119.7	Mo1—O3—H2O	120.8 (18)
C7—C8—C9	120.37 (19)	H1O—O3—H2O	109 (3)
O1—C1—C2—N1	-46.2 (2)	C3—C4—C9—O2	0.6 (3)
N1—C3—C4—C9	8.2 (3)	C5—C4—C9—C8	0.8 (3)
N1—C3—C4—C5	-174.14 (18)	C3—C4—C9—C8	178.44 (16)
C9—C4—C5—C6	-1.7 (3)	C4—C3—N1—C2	-178.44 (18)
C3—C4—C5—C6	-179.50 (19)	C4—C3—N1—Mo1	7.0 (3)
C4—C5—C6—C7	1.2 (3)	C1—C2—N1—C3	-149.68 (18)
C5—C6—C7—C8	0.3 (3)	C1—C2—N1—Mo1	25.66 (19)
C6—C7—C8—C9	-1.3 (3)	C2—C1—O1—Mo1	51.2 (2)
C7—C8—C9—O2	178.67 (17)	C8—C9—O2—Mo1	152.01 (14)
C7—C8—C9—C4	0.7 (3)	C4—C9—O2—Mo1	-30.1 (2)
C5—C4—C9—O2	-177.10 (16)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O3—H1O...O1 <sup>i</sup>	0.73 (2)	1.97 (3)	2.6656 (19)	161 (3)
O3—H2O...O4 <sup>ii</sup>	0.78 (3)	2.07 (3)	2.8425 (19)	173 (3)

Symmetry codes: (i)  $-x+1, -y, -z+1$ ; (ii)  $x, -y+1/2, z-1/2$ .