


 Cite this: *Chem. Commun.*, 2014, 50, 12116

 Received 8th July 2014,
 Accepted 18th August 2014

DOI: 10.1039/c4cc05252h

www.rsc.org/chemcomm

Palladium-catalyzed cyclization of benzamides with arynes: application to the synthesis of phenaglydon and *N*-methylcrinasiadine†


Sandeep Pimparkar and Masilamani Jegannmohan*

***N*-Methyl or methoxy benzamides reacted with benzyne in the presence of Pd(OAc)₂, organic acid and K₂S₂O₈ in CH₃CN yielding tricyclic phenanthridinone derivatives in good yields.**

Phenanthridinones are the key core units found in various natural products and biologically active molecules.¹ These molecules are used as potential PARP-1 inhibitors in anticancer drugs as well as neurotrophin activity enhancers for the treatment of nerve diseases.² Traditionally, phenanthridinones are synthesized by cyclization of nitrocarbonyl-biphenyls, Beckmann/Schmidt rearrangement of fluorenones and photoinduced rearrangement of 2-halobenzamides.³ However, in these reactions, the preparation of key starting materials needs more steps and the overall yields observed are lower. Subsequently, phenanthridinones are prepared by a palladium-catalyzed homo coupling of *ortho*-halo benzamides and coupling of aromatic halides with *ortho*-halo benzamides.⁴ Very recently, Larock's group reported the synthesis of phenanthridinones *via* a palladium-catalyzed cyclization of *ortho*-halo *N*-substituted benzamides with benzyne (eqn (1)).⁵ But a preactivated carbon-halogen partner on the aromatic moiety is required for the reaction. Apart from these reactions, phenanthridinones are prepared by a metal-catalyzed *ortho* arylation of benzamides with iodobenzenes or aromatic boronic acids or electron-rich aromatics followed by intramolecular C–N bond formation.⁶


Transition metal-catalyzed cyclization of heteroatom substituted aromatics with carbon-carbon π -components *via* chelation-assisted C–H bond activation is a practical method to synthesize heterocyclic molecules in one pot.⁷ By using nitrogen containing chelating groups such as amide, oxime and imine substituted aromatics or alkenes, various nitrogen-containing mono- and bicyclic heterocycles are prepared through the consecutive C–C and C–N bond formation.⁷ In the cyclization reaction, alkynes, alkenes and allenes are extensively used as carbon-carbon π -components (eqn (1)). However, benzyne as a π -component has not been well explored in the literature. In fact, there are several challenges to utilize benzyne as a π -component in the reaction due to its high reactivity. It is very important to note that in the cyclization of substituted aromatics with benzyne, a tricyclic ring system can be constructed in one pot.

We focused on utilization of a highly reactive benzyne as a π -component in the cyclization reaction. Initially, we tried the cyclization of *N*-methoxy 4-methoxy benzamide (**1a**) with *o*-(trimethylsilyl)aryl triflate (**2a**) in the presence of [RuCl₂(*p*-cymene)]₂, AgSbF₆ and Cu(OAc)₂ in CH₃CN at 100 °C for 12 h (Table 1, entry 1). CsF in CH₃CN was used to generate benzyne from benzyne precursor **2a**.⁸ In the reaction, only *N*-arylated benzamide **3a** was observed in 30% yield and the expected cyclization product **4a** was not observed. Next, the cyclization reaction of **1a** with **2a** was examined in the presence of Pd(OAc)₂ (5 mol%) and acetic acid or CF₃COOH (10.0 equiv.) in CH₃CN (entries 2 and 3). In the reaction, no *N*-arylation product **3a** or cyclization product **4a** was observed. It seems AcOH or CF₃COOH might quench the CsF base. Then, acetic acid was replaced by the sterically hindered pivalic acid (entry 4). Interestingly, in the reaction, the expected cyclization product **4a** was observed in 35% yield and competitive product **3a** was also observed in 10% yield. To avoid product **3a**, the reaction was tested with a catalytic amount of 1-adamantanecarboxylic acid (Adm-1-COOH) (30 mol%) (entry 5). Surprisingly, in the reaction, product **4a** was observed in 45% yield and no *N*-arylated product **3a** was observed. To increase the yield of product **4a**, the reaction was examined with oxidants (1.0 equiv.) (entries 6–11). Interestingly, using K₂S₂O₈, product **4a** was observed in 74% GC yield and 66%



Department of Chemistry, Indian Institute of Science Education and Research, Pune 411021, India. E-mail: mjegannmohan@iiserpune.ac.in

† Electronic supplementary information (ESI) available: Detailed experimental procedures and spectroscopic data. CCDC 1012984. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4cc05252h


Table 1 Cyclization of benzamide **1a** with benzyne precursor **2a**^a


Entry	Catalyst/additive	Oxidant	3a/4a yield ^b (%)
1	[RuCl ₂ (<i>p</i> -cymene)] ₂ /AgSbF ₆ ^c	Cu(OAc) ₂	30/0
2	Pd(OAc) ₂ /AcOH	—	—
3	Pd(OAc) ₂ /TFA	—	—
4	Pd(OAc) ₂ /pivalic acid	—	10/35
5	Pd(OAc) ₂ /Adm-1-COOH ^d	—	0/45
6	Pd(OAc) ₂ /Adm-1-COOH ^d	Ag ₂ O	0/33
7	Pd(OAc) ₂ /Adm-1-COOH ^d	Ag ₂ CO ₃	0/27
8	Pd(OAc) ₂ /Adm-1-COOH ^d	AgOAc	0/45
9	Pd(OAc) ₂ /Adm-1-COOH ^d	K ₂ S ₂ O ₈	0/74
10	Pd(OAc) ₂ /Adm-1-COOH ^d	PhI(OAc) ₂	—
11	Pd(OAc) ₂ /Adm-1-COOH ^d	(NH ₄) ₂ S ₂ O ₈	—

^a Reactions conditions: **1a** with **2a** (1.5 equiv.) in the presence of cat (5.0 mol%), additive (10.0 equiv.) and oxidant (2.0 equiv.) in CH₃CN at 100 °C for 12 h. ^b GC yield. ^c AgSbF₆ (20 mol%) was used. ^d Adm-1-COOH (30 mol%) was used.



isolated yield (entry 9). Remaining oxidants were partially effective or totally ineffective, yielding **4a** in 0–45% yields (entries 6–11).

The scope of the catalytic reaction was tested with substituted *N*-methoxy benzamides **1b–i** (Scheme 1). 2-Methoxy (**1b**), 4-methyl (**1c**) and *N*-methoxy benzamides (**1d**) underwent cyclization with **2a**, yielding phenanthridinones **4b–d** in 55%, 62% and 61% yields, respectively. Next, the cyclization reaction was tested with unsymmetrical benzamides. *N*-Methoxy 3,4-dimethoxy benzamide (**1e**) and *meta* methoxy benzamide **1f** afforded phenanthridinones **4e** and **4f** in 61% and 58% yields, respectively, in

Scheme 1 Scope of the *N*-methoxy benzamides **1**.


which the *ortho* C–H bond activation takes place selectively at a sterically less hindered side. Whereas, benzamide **1g** provided mixtures of regioselective cyclization products **4g** and **4g'** in 60% combined yield in a 1.5:1 ratio. Furthermore, the cyclization reaction was tested with 4-bromo and 4-chloro benzamides **1h** and **1i**. However, only *N*-arylated benzamides **3h** and **3i** were observed in 47% and 51% yields, respectively, and the expected cyclization products **4** were not observed. Similarly, 4-trifluoromethyl, 4-cyano and 4-nitrobenzamides were also not favorable for the reaction. These results clearly reveal that electron-donating substituents on the aromatic moiety of benzamides favor the *ortho* C–H bond activation–cyclization reaction. But, halogen and electron deficient aromatic benzamides favor only competitive nucleophilic addition of the free N–H moiety of benzamide with benzyne. Surprisingly, the palladacycle of 4-chloro benzamide **5a** reacted with benzyne precursor **2a**, yielding cyclization product **4h** in 45% yield. This result clearly says that the *ortho* C–H bond activation process is very slow in the electron deficient benzamides, and the competitive nucleophilic addition is very fast. Although, at present, the catalytic reaction was compatible with only electron-rich benzamides, it has been shown that a highly reactive benzyne can be used as a π -component for the cyclization reaction.

The cyclization reaction was also examined with benzyne precursors **2b–e** (Scheme 2). Treatment of benzamide **1a** with benzyne precursors **2b** and **2c** gave phenanthridinones **4i** and **4j** in 61% and 55% yields, respectively. Interestingly, synthetically

Scheme 2 Scope of the benzyne precursors **2b–e**.Scheme 3 Scope of the *N*-methyl benzamides **6a–d**.

useful benzyne precursors **2d** and **2e** reacted with **1e**, giving products **4k** and **4l** in 70% and 63% yields, respectively.


The catalytic reaction was further tested with other *N*-methyl benzamides (Scheme 3). *N*-Methyl benzamide (**6a**) underwent cyclization with **2a**, providing *N*-methyl phenanthridinone **7a** in 49% yield, in which *ortho* C–H bond activation takes place selectively at the sterically less hindered side. Further, *N*-methyl 4-methoxy benzamide (**6b**) and *N*-methyl benzamide (**6c**) reacted with **2a** or **2c** affording cyclization products **7b–d** in 45%, 43% and 38% yields, respectively. Treatment of *N*-methyl benzamide **6d** with **1a** gave natural product *N*-methylcrinasiadine^{9b} (**7e**) in 30% yield and other regioisomer **7e'** in 25% yield. It is important to point out that natural product *N*-methylcrinasiadine (**7e**) shows several biological activities.¹




Later, OMe groups on the cyclic amides of **4c** and **4d** were cleaved into the natural product phenaglydon^{9a,b} **8a** in 69% yield and 6(5*H*)-phenanthridinone **8b** in 67% yield under the photochemical irradiation conditions^{7a,b} (eqn (2)). Later, compound **8b** underwent nitration at the C-5 position of phenanthridinone in the presence of HNO₃/H₂SO₄, providing 5-nitrophenanthridinone **9** in 75% yield. It is important to note that compound **9** is a key precursor for the preparation of anticancer drug PJ34.^{9c}

A possible reaction mechanism is proposed in Scheme 4 to account for the present cyclization reaction. Coordination of the amide nitrogen of benzamide **1** to the palladium species followed by *ortho*-metalation provides a five-membered palladacycle intermediate **5**. Coordinative insertion of benzyne **10** into intermediate **5** yields a seven-membered palladacycle intermediate **11**. Subsequent C–N bond formation and reductive elimination afford product **4** and regenerate the active palladium species in the presence of RCOOH and K₂S₂O₈.

Apart from the above proposed mechanism, other possible pathways such as *ortho*-arylation of benzamide with benzyne yielding product **12** followed by intramolecular C–N bond formation or



Scheme 4 Proposed mechanism.



Scheme 5 Mechanistic investigation.

N-H arylation of benzamide with benzyne providing compound **3** followed by intramolecular dehydrogenative aryl–aryl coupling are also possible.⁷ To support the proposed mechanism in Scheme 4, the following reactions were done (Scheme 5). *ortho*-Arylated benzamide **12** was prepared separately and treated with Pd(OAc)₂, CsF and K₂S₂O₈ in CH₃CN at 100 °C for 12 h. In the reaction, no cyclization product **4a** was observed. Subsequently, *N*-arylated benzamide **3a** was treated with Pd(OAc)₂ and K₂S₂O₈ under similar reaction conditions. However, no cyclization product **4a** was observed. Further, a five-membered palladacycle intermediate **5b** was prepared separately and treated with benzyne precursor **2a** in the presence of CsF in CH₃CN at 100 °C for 12 h. As expected, the cyclization product **4f** was observed in 75% yield. These results clearly revealed that the present reaction proceeds *via* a coordinative insertion pathway. To support the hypothesis that benzyne is involved in the cyclization reaction, the reaction of benzamide **1e** with unsymmetrical benzyne precursor **2g** was performed. In the reaction, a mixture of regioisomeric compounds **4n** and **4n'** was observed in 53% combined yield in a 2:1 ratio. The lack of regioselectivity of the reaction is consistent with insertion of unsymmetrical benzyne into a Pd–carbon bond in intermediate **5**.

In conclusion, we have demonstrated a palladium-catalyzed oxidative cyclization of *N*-substituted benzamides with benzyne providing phenanthridinones with diverse substituents.

We thank the DST (SR/S1/OC-26/2011), India for the support of this research. S. P. thanks the BRNS for a fellowship.

Notes and references

- (a) F. Aprile-Garcia, M. Antunica-Noguerol, M. L. Budzinski, A. C. Liberman and E. Arzt, *Endocr. Connect.*, 2014, **3**, 1; (b) S. E. Nasrabady, A. Kuzhandaivel, M. Mladinic and A. Nistri, *Cell. Mol. Neurobiol.*, 2011, **31**, 503; (c) Z. Jin, *Nat. Prod. Rep.*, 2009, **26**, 363; (d) J. P. Michael, *Nat. Prod. Rep.*, 1995, **12**, 465.
- (a) T. Harayama, H. Akamatsu, K. Okamura, T. Miyagoe, T. Akiyama, H. Abe and Y. Takeuchi, *J. Chem. Soc., Perkin Trans. 1*, 2001, 523; (b) T. Harayama, Y. Akiyama, K. Nakano, H. Shibaie, A. Akamatsu, H. Hori, H. Abe and Y. Takeuchi, *Synthesis*, 2002, 237; (c) V. Holl, D. Coelho, D. Weltin, P. Dufour and P. Bischoff, *Anticancer Res.*, 2000, **20**, 3233; (d) S. Patil, S. Kamath, T. Sanchez, N. Neamati, R. F. Schinazi and J. K. Buolamwini, *Bioorg. Med. Chem.*, 2007, **15**, 1212.
- (a) A. K. Mohanakrishnan and P. C. Srinivasan, *J. Org. Chem.*, 1995, **60**, 1939; (b) R. Kleppinger, C. P. Lillya and C. Yang, *J. Am. Chem. Soc.*, 1997, **119**, 4097; (c) M. G. Banwell, D. W. Lupton, X. Ma, J. Renner and M. O. Sydnes, *Org. Lett.*, 2004, **6**, 2741; (d) R. Kleppinger, C. P. Lillya and C. Yang, *J. Am. Chem. Soc.*, 1997, **119**, 4097; (e) L. M. N. Meseroll, J. R. McKee and M. Zanger, *Synth. Commun.*, 2011, **41**, 2557;

- (f) A. Padwa, M. Dimitroff, A. G. Waterson and T. Wu, *J. Org. Chem.*, 1998, **63**, 3986; (g) H. Suzuki, T. Enya and Y. Hisamatsu, *Synthesis*, 1997, 1273.
- 4 (a) T. Furuta, Y. Kitamura, A. Hashimoto, S. Fujii, K. Tanaka and T. Kan, *Org. Lett.*, 2007, **9**, 183; (b) P. H. Bernardo, W. Fitriyanto and C. L. L. Chai, *Synlett*, 2007, 1935; (c) R. Ferraccioli, D. Carenzi, O. Rombola and M. Catellani, *Org. Lett.*, 2004, **6**, 4759; (d) V. Rajeshkumar, T.-H. Lee and S.-C. Chuang, *Org. Lett.*, 2013, **15**, 1468.
- 5 C. Lu, A. V. Dubrovskiy and R. C. Larock, *J. Org. Chem.*, 2012, **77**, 8648, references therein.
- 6 (a) P. Thansandote and M. Lautens, *Chem. – Eur. J.*, 2009, **15**, 5874; (b) T. Satoh and M. Miura, *Chem. – Eur. J.*, 2010, **16**, 11212; (c) G. Song, F. Wang and X. Li, *Chem. Soc. Rev.*, 2012, **41**, 3651; (d) H. Wang and F. Glorius, *Angew. Chem., Int. Ed.*, 2012, **51**, 7318; (e) P. B. Arokiam, C. Bruneau and P. H. Dixneuf, *Chem. Rev.*, 2012, **112**, 5879; (f) L. Ackermann, *Acc. Chem. Res.*, 2014, **47**, 281; (g) T. W. Lyons and M. S. Sanford, *Chem. Rev.*, 2010, **110**, 1147.
- 7 (a) T.-T. Yuan, D.-D. Li and G.-W. Wang, *Angew. Chem., Int. Ed.*, 2011, **50**, 1380; (b) J. Karthikeyan and C.-H. Cheng, *Angew. Chem., Int. Ed.*, 2011, **50**, 9880; (c) J. Karthikeyan, R. Haridhadharan and C.-H. Cheng, *Angew. Chem., Int. Ed.*, 2012, **51**, 12343; (d) N. Senthilkumar, K. Parthasarathy, G. Parthasarathy and C.-H. Cheng, *Chem. – Asian J.*, 2013, **8**, 2175; (e) C. S. Yeung, X. Zhao, N. Borduas and V. M. Dong, *Chem. Sci.*, 2010, **1**, 331.
- 8 Y. Himeshima, T. Sonoda and H. Kobayashi, *Chem. Lett.*, 1983, 1211.
- 9 (a) G. Wurz, O. Hofer and H. Greger, *Nat. Prod. Lett.*, 1993, **3**, 177; (b) R. Suau, A. I. Gomez and R. Rico, *Phytochemistry*, 1990, **29**, 1710; (c) Z. Tu, W. Chu, J. Zhang, C. S. Dence, M. J. Wenck and R. H. Mach, *Nucl. Med. Biol.*, 2005, **32**, 437.